DIRECTIONS FOR USE


In hips doct is toxic to birds and only welflife 150 mid. I stanguage any body. I willer with this an today emply only as specification this label. In abote of left over material or empty conthe ray by burning or burying in non-grop land. addy from water supplies. Stop out of smoke, Fig. at a use container, CARZOL SP a modor his time to honey bees. Aveid many where bond are active. Do not allow animals to graze in christiand areas treated with CARZOL SP

Purchase or use of this product shall constitute acceptance of the following terms of sale. NOR-AM Agricultural Products, Inc. (NOR AM) warrants that this product conforms to the chemical descriptions on this label and is reasonably fit for the purposes specified hereon. By way of limitation but not in derogation of the directions and statements printed on this label. Seller makes no warranty and no representation and NOR-AM makes no further warranty or representation, expressed or implied, including but not limited to any warranty as to MERCHANTABILITY, and no agent of NOR AM is authorized to do so except in writing, with a specific reference to the foregoing warranty. Since neither Seller nor NOR-AM has any control over the handling, storage, use or conditions of use of this product, Buyer er user assumes all risks and liability arising from handling, storage or use of this product under


abnormal conditions or not in strict compliance with the directions and precautionary.

IMPORTANT:

PEAD BEFORE PURCHASE OR USE



Decemed by and Trademark* of behenng A. G. Berlin, Bergkamen 🙀



MITICIDE/INSECTICIDE

Packed In Water Soluble Bags To Be Sold As One Unit Only

STORE IN A DRY PLACE AT A TEMPERATURE BELOW 125°F

KEEP OUT OF REACH OF CHILDREN

Not for use or storage in or around the home. Call a physician immediately in case of exposure.

This product is poisonous if swallowed. Do not get in eyes, on skin or on clothing. Do not breathe dust or spray mist. Do not contaminate food and feedstuffs. Wear protective clothing, goggles and rubber (natural or synthetic) gloves when handling this material. Wash before eating or smoking. Wash contaminated clothing before re-use.

If exposure to dust or spray mist cannot be avoided wear a self-contained breathing apparatus.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY

If swallowed-give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician immediately. If on skin-wash immediately with soap and water, If in eyesflush with water for at least 15 minutes and get medical attention.

NOTICE TO PHYSICIAN:

CARZOL SP is a carbamate which acts as a cholinesterase inhibitor.

Specific antidote—Atropine.

Do not use 2 PAM

See Side Panel For Additional Precautions and Injections for the

No's #396196-1-3542850 -

NET CONT.: 10 X 2½ EBS.

A S MICKEN ACTURED FOR NOR-AM.

AGRICULTURAL PRODUCTS, MC

CHICAGO, ILLINOIS BORGE

FPA Est. No. 49967 NEED

Unaffection U.S.A.

This package contains ten water soluble bags of Carzol SP each one containing 255 paged of product. Carzol SP is completely soluble in water. Preparation of a pre-mix or through adding to the spray tank is not necessary. Add the required quaker of soluble back without opening them, if recly to the spray tank whole the time is to say trivial and inrunning. Make sure that the bags are completely displayed in the tack provide adding to.

PRECAUTIONS

Do not handle bags with moist hands.

To protect unused bags from moisture, immediately ciose the outside bag after soluble bags and place the box in a dry place.

Do not expess bars to surlight, next or moreture.

MOTICE: CAR/OI SP in and Mahie on a Paline water. Do not propose the option models of just before application. Do not prepare more spring maximo thin will be used within 4 his

CROP	PEST	APPLICATION
ORANGES, LEMONS AND TANGERINES (California & Arizona only)		Use 8 oz. CARZOL SP per 100 gallons of water Apply for outside coverage in 100 to 300 gallons of water per acre; the to 132 lbs. CARZOL SP per acre) depending on the size of the trees. Repeat application as needed.
	RED MITE	Use 8 oz. CARZOL SP per 100 gallons of water. Apply in a full coverage spray in 700 to 1000 gallons of water per acre (3½ to 5 lbs. CARZOL SP per acre) depending on the size of the frees.
ORANGES, LEMONS TANGERINES AND GRAPEFRUIT (in Florida)	RUST MITE	Use 1 to 2 oz. CARZOL SP per 100 gallons of water, apply in full coverage spray. Use 7 to 20 oz. CARZOL SP per acre depending on the size of the irres. Repeat application as needed.
ORANGES, LEMONS TANGERINES AND GRAPEFRUIT (in Texas)	RUST MITE	Use 2 to 4 oz. CARZOL SP per 100 gallons of water, apply in full coverage spray. Use 10 to 20 oz. CARZOL SP per acre depending on the size of the trees. Repeat application as needed.
APPLES AND PEARS	EUROPEAN RED MITE, TWO-SPOTTED MITE	Use 4 to 8 oz. CARZOL SP per 100 gallons of water Apply m full coverage sprays in 400 to 800 gallons of water per acre (1 to 4 lbs. of CARZOL SP per acre) depending on the size of the trees and severity of the mite infestation. Repeat application as needed.
PEACHES AND NECTARINES (West Coast only)	TWO SPOTTED MITE EUROPEAN RED MITE MCDANIEL MITE	Apply 4 oz. CARZOL SP per 100 gallons of spray (1 to 1¼ lbs. CARZOL SP per acre) when mites appear. Repeat as necessary.
NECTARINES (California only)	THRIPS .	Apply 8 oz. CARZOL SP per 100 gailons of spray (2 lbs. CARZOL SP per acre) when thrips infest blossoms through petal fall. Apply at night or while bees are not working in the blooms.
PLUMS, PRUNES (West Coast only)	TWO SPOTTED MITE EUROPEAN RED MITE	Apply 4 oz. CARZOL SP per 100 gallons of spray (1 to 1½ lbs. CARZOL SP per acre) when mites appear. Repeat as necessary.
ALFALFA (Seed crop only, in Calif. Oregon, Washington, Idaho and Nevada. Use pre-bloom only on alfalfa in all areas except Califorma.)	LYGUS BUGS	Use ½ to 1 lbs. CARZOL SP per acre. Apply when lygus bugs reach damaging numbers. Repeat application as needed. Consult your State Agricultural Experiment Station or State Extension Service Specialists for proper timing.
	CONSPERSE STINK BUGS (California only)	Use 1 lb. CARZOL SP per acre for knockdown of heavy nymphal population. A lygus control program will control stink bugs.
		When applied by air on alfalfa, use a minimum of 5 gallons per acre. When applied by ground use a minimum of 20 gallons per acre.

NOTE

To avoid excessive residues, observe the following limitations: CITRUS: Do not apply more than 5 lbs. CARZOL SP per acre new 12 month period. Allow 30 days between applications and 7 days before harvest. APPLES PEARS. No not apply more than 4 Hrs. CARZOL SE per acce in one crop season. Do not apply less than 7 days before harvest. PLACHES, to not apply more than 2.5 lbs. CARZOL SP per acre in one crop season. Do not apply less than 21 days before harvest. NECLARINES. Do not apply more than 4 lbs. per acre in one crop season. Do not apply less than 14 days before harves! PLUMS, PRUNES: Do not apply more than 3 ibs per acre in one crop season. Do not apply less than 7 days before harvest. ALFALFA (seed crop only) lise this product only on pure stands of affalfa to be harvested for seed. Do not allow spray to drift onto adjacent grops. Do not spray while bees are foraging. Do not allow livestock to graze heated helds. Do not use treated crops for live stock feed. Crop residues are not to be fed to livestock or baled for such use. Fields treated with CARZOL SP must be cleared after harvest by burning or by disking into the soil. CARZOL SP does not control aghids. If aphids appear add an aphicide to the spray.