


PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

2139-99

06/13/1975

CARZOL SP, EPA Reg. No. 2139-99

Amended Label Claim

Citrus (Aerial Application for Thrips Control)

DIRECTIONS FOR USE

CARZOL SP is completely soluble in water. Preparation of a premix or slurry before adding to the spray tank is not necessary. Add the desired amount of CARZOL directly to the spray tank 1/2 full of water with the agitator running. Then add the remainder of the water.

NOTICE: CARZOL SP is not stable in alkaline water. Do not prepare the spray mixture until just before application. Do not prepare more spray mixture than will be used within 4 hours.

CROP	PEST	APPLICATION
Oranges, lemons, tangerines, tangelos, grapefruit (California and Arizona)	Thrips	Use 8 oz. CARZOL SP per 100 gallons of water when thrips appear. Apply for outside coverage in 100 to 300 gallons of water per acre (1/2 to 1-1/2 lbs. CARZOL SP per acre) depending on the size of the trees. For aerial application apply 1 to 1-1/2 lbs. CARZOL SP per acre in a minimum of 10 gallons of water. Repeat application as needed.

NOTE

To avoid excessive residues, observe the following limitations:
CITRUS: Do not apply more than 5 pounds CARZOL SP per acre in a 12-month period. Allow 30 days between applications and 7 days before harvest.

ML:slt
3/5/75

NEXT

LABEL

6-13-1975


ML:slt
3/5/75

NEXT

LABEL

2139-99

6-13-1975



CARZOL SP, EPA Reg. No. 2139-99

Amended Label Claim

Control of Twospotted Mite in Seed Alfalfa

DIRECTIONS FOR USE

CARZOL SP is completely soluble in water. Preparation of a premix or slurry before adding to the spray tank is not necessary. Add the desired amount of CARZOL SP directly to the spray tank 1/2 full of water with the agitator running. Then add the remainder of the water.

NOTICE: CARZOL SP is not stable in alkaline water. Do not prepare the spray mixture until just before application. Do not prepare more spray mixture than will be used within 4 hours.

CROP	PEST	APPLICATION
ALFALFA (Seed crop only, in Calif., Oregon, Washington, Idaho and Nevada). Use pre-bloom only on alfalfa in all areas except California.	Twospotted Mite	Use 1/2 to 1 pound CARZOL SP per acre. Apply when mites appear. When applied by air use a minimum of 5 gallons of spray per acre. When applied by ground use a minimum of 20 gallons per acre. Repeat application as needed.

NOTE

To avoid excessive residues, observe the following limitations. ALFALFA (seed crop only): Use this product only on pure stands of alfalfa to be harvested for seed. Do not allow spray to drift onto adjacent crops. Do not spray while bees are foraging. Do not allow livestock to graze treated fields. Do not use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. Fields treated with CARZOL SP must be cleared after harvest by burning or by disking into the soil. CARZOL SP does not control aphids. If aphids appear add an aphicide to the spray.

ML:slt
3/5/75

NEXT

LABEL

2139-99

6-13-1975

1/1

CARZOL SP, EPA Reg. No. 2139-99

Amended Label Claim


Peaches, Nectarines, Plums, Prunes

DIRECTIONS FOR USE

CARZOL SP is completely soluble in water. Preparation of a premix or slurry before adding to the spray tank is not necessary. Add the desired amount of CARZOL SP directly to the spray tank 1/2 full of water with the agitator running. Then add the remainder of the water.

NOTICE: CARZOL SP is not stable in alkaline water. Do not prepare the spray mixture until just before application. Do not prepare more spray mixture than will be used within 4 hours.

CROP	PEST	APPLICATION
Peaches, Nectarines (West Coast only)	Twospotted Mite European Red Mite McDaniel Mite	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. Apply by ground in full cover dilute spray at 400-500 gallons per acre. For concentrate spray increase the concentration so that the amount of CARZOL SP is 1 to $1\frac{1}{4}$ lbs./acre in at least 50 gallons of spray per acre. Repeat application as needed.
Plums, Prunes (West Coast only)	Twospotted Mite European Red Mite McDaniel Mite	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. Apply by ground in full cover dilute spray at 400-600 gallons per acre. For concentrate spray increase the concentration so that the amount of CARZOL SP is 1 to $1\frac{1}{2}$ lbs./acre in at least 50 gallons of spray per acre. Repeat application as needed.



NOTE

To avoid excessive residues, observe the following limitations:
 PEACHES: Do not apply more than 2 1/2 lbs. CARZOL SP per acre in one crop season. Do not apply less than 21 days before harvest. NECTARINES: Do not apply more than 4 lbs. per acre in one crop season. Do not apply less than 14 days before harvest. PLUMS, PRUNES: Do not apply more than 3 lbs. per acre in one crop season. Do not apply less than 7 days before harvest.

ML:slt
3/5/75