



Lanex PLUS

COTTON HERBICIDE

For post-emergence weed control in cotton

THIS MIXTURE CONTAINS 2.0 LB. LANEX 80 WP AND 3.3 LB. MSMA PER GAL.

E.P.A. Registration No. 2139-107

ACTIVE INGREDIENT

Fluometuron* 13.2%
Monosodium acid methanearsonate** 27.6%

INERT INGREDIENTS 59.2%
TOTAL 100.0%

*1, 1-dimethyl-3-(4-chloro-2-trifluoro-m-tolyl) urea

**Total arsenic, as elemental, all in water soluble form, 12.7%

CAUTION: Keep out of reach of children

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands with soap and water before eating, drinking, or smoking. Wear clean clothing. Do not store near food or feed. Destroy empty container. Do not graze or feed plants from treated areas or gin trash to livestock. Do not contaminate water used for irrigation or domestic purposes.

NOTE: Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

IMPORTANT: Read the entire directions for use and the non-warranty on the left and right panels before using this product. See left and right panels for additional precautions.

U.S. PATENT NO. 3,134,665

NET CONTENTS: 1 GAL.



NOR-AM

AGRICULTURAL PRODUCTS, INC.

ACCEPTED

Sept 20, 1977

U.S. PATENT NO. 3,134,665
E.P.A. REG. NO. 2139-107
2139-107

GENERAL INFORMATION: LANEX PLUS is effective for post-emergence control of broad-leaf and grass weeds in cotton, such as

Grass Weeds

Barnyardgrass	Crowfootgrass	Fall Panicum	Nutgrass
Bracharia	Dallisgrass	Foxtail	Ryegrass
Crabgrass	Goosegrass	Johnsongrass	Sandbur

Broadleaf Weeds

Buttonweed	Jimsonweed	Prickly sida	Ragweed
Cocklebur	Lambsquarters	(ironweed or teaweed)	Sesbania
Florida pussley	Morningglory	Puncturevine	Sicklepod
Goathead	Pigweed	Purslane	Smartweed
			Tumbleweed


West Texas, Western Oklahoma and Northeast New Mexico: Do not use this product on sandy, loamy sand, or fine sandy loam soils.

NOTE: DO NOT STORE AT TEMPERATURES BELOW FREEZING KEEP CONTAINER CLOSED

APPLICATION DIRECTIONS:

NOTE: Shake well before using. Do not make more than two applications of LANEX PLUS to the same crop or field in any one year. Apply only when cotton is 3 inches high to first bloom. Apply as a directed spray, avoiding cotton foliage. Keep children and livestock off treated areas until this material has been washed into the soil. Do not graze treated field or feed plants or gin trash from treated areas to livestock. This product contains surfactant. Do not add surfactant or damage may result. Do not plant crops other than cotton within six months after last application

Apply LANEX PLUS at the broadcast rate of 1/2 gallon per acre, in 25 to 40 gallons of water per acre. Direct spray to the base of the cotton plants, avoiding the cotton foliage. Best results obtained when applied to young, actively growing weeds at air temperatures above 70°F



For band applications, the following formula may be used to calculate the amount of LANEX PLUS required:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times 0.5 = \text{amount, in gallons needed for band treatments}$$

Slight burning and a reddish discoloration of the cotton leaf may occur occasionally following treatment with products containing MSMA, but the cotton plants will develop normally.

Equipment: Use conventional spray equipment and Tee Jet nozzle tips 8002 to 8006, or equivalent. Screen size should be 50 mesh.

MIXING INSTRUCTIONS: LANEX PLUS is a highly concentrated formulation and should be thoroughly shaken before mixing. Pour recommended amount of LANEX PLUS into nearly filled spray tank, or premix in a bucket half filled with water and pour mixture into nearly filled spray tank. By-pass agitation should terminate at the bottom of the tank to minimize foaming. Avoid leaving spray mixture in tank without constant agitation.

DO NOT ADD SURFACTANT TO THE MIX.

After use, clean equipment thoroughly. Do not contaminate water by dumping unused spray or washings in ponds or streams. Destroy empty containers.

IMPORTANT: READ BEFORE PURCHASE OR USE

Purchase or use of this product shall constitute acceptance of the following terms of sale. NOR-AM Agricultural Products, Inc. (NOR-AM) warrants that this product conforms to the chemical descriptions on this label and is reasonably fit for the purposes specified hereon. By way of limitation but not in derogation of the directions and statements printed on this label, Seller makes no warranty and no representation and NOR-AM makes no further warranty or representation, expressed or implied, including but not limited to any warranty as to MERCHANTABILITY, and no agent of NOR-AM is authorized to do so except in writing with a specific reference to the foregoing warranty. Since neither Seller nor NOR-AM has any control over the handling, storage, use or conditions of use of this product, Buyer or user assumes all risks and liability arising from handling, storage or use of this product under abnormal conditions or not in strict compliance with the directions and precautionary statements hereon. All claims related to the purchase, handling, storage or use of this product shall be barred unless written notice is given to NOR-AM within 30 days after the loss is discovered. Liability of the Seller or NOR-AM for any damages arising from a breach of this warranty shall be limited to exclude consequential commercial damages.

725

DO NOT RE-USE THIS CONTAINER

PRINTED IN U.S.A.