



Center Panel

RECEIVED

JUN 30 1983

2139-124



PREVEX<sup>TM</sup>

FUNGICIDE FOR CONTROL OF PYTHIUM AND PHYTOPHTHORA IN ORNAMENTALS

EPA Reg. No. 2139-

Active ingredient: Propamocarb hydrochloride*	66.5%
*Propyl [3-(dimethylamino)propyl]carbamate monohydrochloride	
Inert ingredients	33.5%
Total	100%

Contains 6.0 lbs. active ingredient per gallon.

**KEEP OUT OF THE REACH OF CHILDREN**  
**CAUTION!**

**STATEMENT OF PRACTICAL TREATMENT**  
If swallowed: administer gastric lavage. If in eyes: immediately flush with plenty of water.  
If on skin: wash with soap and water. Get medical attention if irritation persists.

**HAZARDS TO HUMANS**  
May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Do not enter treated areas until spray deposit has dried.

**ENVIRONMENTAL HAZARDS**  
Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning equipment or disposals of wastes.

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS

Net Contents: 1 Quart




MANUFACTURED FOR  
**NOR-AM.**  
AGRICULTURAL PRODUCTS, INC.  
MARTINVILLE, ILLINOIS 60566

Product of West Germany

BEST DOCUMENT AVAILABLE

Proposed Amendment 7/8/81



Right Panel

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Storage: Keep away from food and feedstuffs. Store in original container, tightly closed.  
Disposal: Do not reuse container. Rinse empty container and bury it and product that cannot be used in a safe place away from water supplies.

Seeding: Use 3 fl. oz. PREVEX/10 gallons of water. Apply to the seeding area using 3 quarts of solution/10 square feet. Treat before sowing of seeds requiring light and after covering of seeds requiring dark for germination.

Transplanting (Seedlings): Use 3 fl. oz. PREVEX/10 gallons of water. Apply to the transplanting area using 3 quarts of solution/10 square feet before transplanting.

Transplanting (Cuttings): Use 2 fl. oz. PREVEX/10 gallons of water. Drench the cutting bed using 2 gallons of solution/10 square feet before setting.

Potting: Use 2 fl. oz. PREVEX/10 gallons water. Dip bare root plants in the solution before potting. After potting drench each pot at the rate of 3.5 fl. oz. solution/4 inch pot. Repeat the drench treatment at 3 to 6 week intervals.

For control of Phytophthora spp on woody ornamentals such as azalea, use 2½ fl. oz. PREVEX/10 gallons of water. Drench each container at the rate of 5 quarts solution/10 square feet of surface. Repeat the drench treatment at 3 to 4 month intervals.

GENERAL INFORMATION

Left Panel

**IMPORTANT:  
READ BEFORE PURCHASE OR USE**

Purchase or use of this product shall constitute acceptance of the following terms of sale. NOR-AM Agricultural Products, Inc. (NOR-AM) warrants that this product conforms to the chemical descriptions on this label and is reasonably fit for the purposes specified hereon. By way of limitation but not in derogation of the directions and statements printed on this label, Seller makes no warranty and no representation and NOR-AM makes no further warranty or representation expressed or implied, including but not limited to any warranty as to MERCHANTABILITY, and no agent of NOR-AM is authorized to do so except in writing, with a specific reference to the foregoing warranty. All claims related to the purchase, handling, storage or use of this product shall be barred unless written notice is given to NOR-AM within 30 days after the loss is discovered. Liability of the Seller or NOR-AM for any damages arising from a breach of this warranty shall be limited to exclude consequential commercial damages.

PREVEX is recommended for prevention of root rot and damping off caused by Pythium spp and Phytophthora spp at all stages of ornamental propagation. For control of other soil borne diseases such as Rhizoctonia spp or Fusarium spp PREVEX can be tank mixed with benomyl or thiophanate-methyl. Use the full rate of each fungicide and follow all directions and restrictions appearing on the labels for these products.

PREVEX is completely water miscible and requires no agitation after initial mixing. Stock solutions of PREVEX should be used within one week of mixing.

PREVEX does not cause phytotoxic symptoms on the following plants when used as directed. Trials on other species should be conducted before extensive use.

African violet	Geranium
Ageratum	Gloxinia
Alyssum	Impatiens
Aster	Lobelia
Azalea	Marigold
Browallia	Pansy
Carnation	Petunia
Celosia	Poinsettia
Chrysanthemum	Portulaca
Coleus	Salvia
Dahlia	Snapdragon
Dusty Miller	Verbena
Easter Lilly	Violet
Ferns	Vinca