

## DIRECTIONS FOR USE

### GENERAL INFORMATION:

St. Augustine Broadleaf Weed Killer is specifically formulated for maximum safety to seedling grasses and 2, 4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill, the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur.

To insure weeds are growing and able to absorb the herbicide, irrigate the lawn 2 or 3 days prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70 degrees F, sprays can be applied at any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dichondra and most ornamental and vegetable crops.

Amount of water in spray is not important except to evenly distribute the chemical. If directions call for 16 oz. in 2 gallons on 1000 square feet, it may be more convenient to use 16 oz. in 4 gallons on 1000 square feet, depending on the sprayer. Generally large volumes of water result in more uniform coverage.

### HOW TO MIX:

- For use on St. Augustine grass, bentgrass and buffalo grass, mix 16 oz. in 2 to 4 gallons of water and uniformly spray 1000 square feet, for example, an area 10 feet wide and 100 feet long. If necessary, repeat treatment in 2 to 3 weeks, if hard to kill weeds persist.
- For use on most other grasses such as bermuda, Bahia, bluegrass, centipede, fescue and rye, mix 20 oz. in 2 to 4 gallons of water and spray uniformly over 1000 square feet. On hard to kill species, or under poor growing conditions, or on older weeds, mix 32 oz. in 2 to 4 gallons of water and spray on 1000 square feet with special attention to thorough coverage of the hard to kill weeds.



## St. Augustine Grass Broadleaf Herbicide

Contains Trimec® Herbicide (1)

**KILLS - DANDELIONS-KNOTWEED-HENBIT-CHICKWEED-PLAINTAINS-SPURGE- and many other broadleaf weeds.**

**ONE QUART COVERS UP TO 2,000 SQUARE FEET  
KEEP FROM FREEZING**

### ACTIVE INGREDIENTS:

|                                                                     |       |
|---------------------------------------------------------------------|-------|
| *Dimethylamine salt of 2, 4-dichlorophenoxyacetic acid              | 0.63% |
| **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid | 2.08% |
| ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)       | 0.25% |

### INERT INGREDIENTS:

|              |                |
|--------------|----------------|
|              | 97.04%         |
| <b>TOTAL</b> | <b>100.00%</b> |

\*2,4-dichlorophenoxyacetic acid equivalent 0.044 pounds per gallon or 0.528%.  
\*\*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.145 pounds per gallon or 1.720%.  
\*\*\*3,6-dichloro-o-anisic acid equivalent 0.018 pounds per gallon or 0.213%.

(1) Trimec® is a registered trademark of PBI-Gordon Corporation, U.S. Patent No. 3,284,186.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS.

Sold by

**I. SCHNEIDER, INC.**  
1429 FAIRMONT AVE., N. W.  
ATLANTA 18, GEORGIA

ACCEPTED  
MAY 14, 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
UNDER NO. 2155-62

### WEEDS CONTROLLED:

icide is especially effective on  
**SHEEP SORREL, KNOTWEED, CHICKWEED, OXALIS** and  
these, it kills virtually all  
which are listed below.

|              |                 |
|--------------|-----------------|
| Bedstraw     | Gallinweed      |
| Black Medick | Ground Ivy      |
| Buckhorn     | Henbit          |
| Bull Thistle | Henbit          |
| Burdock      | Henbit          |
| Buttercup    | Knotted         |
| Chicory      | Lamb's Quarters |
| Chickweed    | Leopards        |
| Clover       | Mallow          |
| Cocklebur    | Morning Glory   |
| Dandelion    | Mustard         |
| Dock         | Oxalis          |

**CAUTION:** Harmful if swallowed, skin, eyes or clothing.

If contact with eyes or clothing.

Avoid drift of spray on ornamental plants, shrubs, plants. Do not contaminate. Do not store in contact with pesticides or fungicides. this use. Coarse sprays are apply to areas underlain by shrubs.

When container is empty, destroy. Never re-use. Destroy and burying in a safe place.

**NOTICE:** Seller warrants the ingredient statement on the label such as weather, compatibility, dilution of application or claims other than those stated.

**CONTROL NUMBER**

E. P. A. Reg. No. 2155-62



# St. Augustine Grass Broadleaf Herbicide

Contains Trimec® Herbicide<sup>(1)</sup>

**LLS - DANDELIONS-KNOTWEED-HENBIT-CHICKWEED-PLAINTAINS-SPURGE-**  
and many other broadleaf weeds.

**ONE QUART COVERS UP TO 2,000 SQUARE FEET**  
**KEEP FROM FREEZING**

## ACTIVE INGREDIENTS:

\*Dimethylamine salt of 2, 4-dichlorophenoxyacetic acid ..... 0.63%  
\*\*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid..... 2.08%  
\*\*\*Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) ..... 0.25%

## INERT INGREDIENTS:

**TOTAL 100.00%**

\*2,4-dichlorophenoxyacetic acid equivalent 0.044 pounds per gallon or 0.528%  
\*\*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.145 pounds per gallon or 1.710%  
\*\*\*3,6-dichloro-o-anisic acid equivalent 0.018 pounds per gallon or 0.213%

(1) Trimec® is a registered trademark of PBI-Gordon Corporation, U.S. Patent No. 3,284,186.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
**SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS.**

Sold by

**I. SCHNEIDER, INC.**  
**1429 FAIRMONT AVE., N. W.**  
**ATLANTA 18, GEORGIA**

**ACCEPTED**

**WEEDS CONTROLLED.** St. Augustine Grass Herbicide is especially effective against weeds such as **SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT.** In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

|              |               |                  |              |
|--------------|---------------|------------------|--------------|
| Bedstraw     | Galinsoga     | Peppergrass      | Sourgrass    |
| Black Medic  | Ground Ivy    | Pigweed          | Velvet Leaf  |
| Buckhorn     | Hawweed       | Plaintains       | Wild Carrot  |
| Bull Thistle | Hogweed       | Poison Ivy       | Wild Garlic  |
| Burdock      | Henbit        | Poison Oak       | Wild Lettuce |
| Buttercup    | Knotweed      | Ragweed          | Wild Onion   |
| Chicory      | Lambsquarter  | Sheep Sorrel     | Yarrow       |
| Chickweed    | Lespedeza     | Shepherd's Purse |              |
| Clover       | Mallow        | Speedwell        | And many     |
| Cocklebur    | Morning Glory | Tansy Mustard    | other broad- |
| Dandelion    | Mustard       | Thistle          | leaf weeds   |
| Dock         | Oxalis        |                  |              |

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. If contact with eyes occurs flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not store in contact with fertilizers, seeds, insecticides or fungicides. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.

**CONTROL**  
**NUMBER**

E. P. A. Reg. No. 2165-62

**CONT.**  
**GALS.**

15A

UNDER THE FEDERAL INSECTICIDE ACT

NOV. 11, 1972