

[2155-39]

**DIRECTIONS FOR USE**

**GENERAL:** For most effective results against the active stages of flying, running and crawling insects, use in a good sprayer adjusted to deliver a fine mist. Shut off ventilating systems. Cover food, food products and utensils. Treat infested areas thoroughly and whenever possible, keep doors and windows closed for 10 minutes after spraying. Spray as often as necessary for control.

**FLYING INSECTS:** For control of flies, mosquitoes, gnats, wasps, hornets, sand flies and certain other flying pests, direct the spray toward the upper portions of the room and saturate the atmosphere with the mist, using 1 ounce to each 3,000 cubic feet of space. Rapidity of initial kill may be accelerated by increasing dosage; however, not more than 1 gallon should be atomized in each 40,000 cubic feet.

**CRAWLING INSECTS:** To control roaches, silverfish, crickets, spiders, spray infested areas thoroughly and direct the liquid behind baseboards, moldings, cabinets, sinks, and window frames, and pipes, toward drains and dark corners, and into crevices, recesses and similar hiding places.

**AGAINST ANT INFESTATIONS,** treat shelves, cabinets and storage areas; spray ant runs, trails, baseboards, thresholds and other places of entry, and treat nests liberally. Repeat treatment when ants are seen.

**IN CONTROLLING FLEAS AND TICKS IN HOUSES,** direct the application toward the lower wall portions, baseboards, rugs, carpets and floors of infested areas. Animal quarters, kennels and bedding should also be sprayed. **CAUTION: DO NOT SPRAY ANIMAL PETS.**

# SANO-CIDE

## DEODORIZED INSECTICIDE CONCENTRATE

**EXCELLENT FOR USE IN FOOD HANDLING AND FOOD PROCESSING ESTABLISHMENTS**

ACTIVE INGREDIENTS 100%

- Petroleum Distillate
- N-Octyl Bicycloheptene Dicarboximide
- \*Technical Piperonyl Butoxide
- Pyrethrins

\*Consists of (butylcarbityl) (6-propylpiperonyl) ether and related compounds

**KILLS BY CONTACT EXPOSED STAGES OF:**

- |                   |                   |                       |
|-------------------|-------------------|-----------------------|
| <b>ROACHES</b>    | <b>MOSQUITOES</b> | <b>CLOTHES MOTHS</b>  |
| <b>SILVERFISH</b> | <b>SKIPPERS</b>   | <b>BEDBUGS</b>        |
| <b>CRICKETS</b>   | <b>GNATS</b>      | <b>CARPET BEETLES</b> |
| <b>SPIDERS</b>    | <b>WASPS</b>      | <b>FLEAS</b>          |
| <b>ANTS</b>       | <b>HORNEYS</b>    | <b>TICKS</b>          |
| <b>FLIES</b>      | <b>SAND FLIES</b> | <b>GRAIN INSECTS</b>  |

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

SEE RIGHT PANEL FOR OTHER PRECAUTIONS

**DIRECTIONS FOR USE**

*(Continued from Left Panel)*

Treat walls, baseboards, floors, closets, furniture, with particular attention to hiding places.

**CLOTHES MOTHS AND CARPET BEETLES:** Remove infested articles from storage, brush thoroughly and air several hours in direct sunlight. Apply spray liberally in closets, bureaus and storage areas directing the liquid into joints and crevices. After airing, infested articles must be treated thoroughly by holding the spray nozzle at least 3 feet away from the fabric and applying the liquid lightly on the entire article. Treat both sides of rugs, carpets and pads, and all exposed and accessible surfaces of overstuffed furniture. Repeat treatment monthly for susceptible articles not stored in moth-tight containers.

**FLOUR BEETLES, WEEVILS, CADELLES, GRAIN BEETLES, MEAL WORMS, SPIDER BEETLES, MEAL MOTHS, SKIPPERS, GRAIN AND CHEESE MITES:** To avoid sources of reinfestation, wormy flour, grain and cereals should be destroyed before treatment. Spray thoroughly all surfaces of shelves, cupboards, pantries, and similar storages, with particular attention to crevices and other hiding places. Repeat treatment as needed.

**CAUTION**

KEEP OUT OF REACH OF CHILDREN  
Do Not Spray On Or Near Open Flame.  
Harmful If Swallowed.

Avoid contamination of food. Use as a space spray in food processing plants only after plant has shut down, food is removed and processing surfaces have been cleaned and covered.

ASCEP  
10/12/67  
2155-39