

DIRECTIONS FOR USE

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns, wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seeds or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results, mow before treating. Do not water lawn within 24 hours after treatment. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot Spray" weeds only. Clover present be damaged if sprayed. Repeat if necessary for hard to kill weeds.


HOW MUCH TO USE -- ENTIRE LAWN

HOSE-END SPRAYER. Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example, to spray 2,000 square feet, measure 2 pints (32 oz.) in jar and add water to the 10 gallon level and mix. Turn on hose and spray as evenly as possible over 2,000 square feet.

Col. 1 Amt. Lawn Weed Killer to Use:		Col. 2 Fill Bottle To: (Gallons)	Col. 3 Area to Treat: (Sq. Feet)
Oz.	Pint		
16	1	5	1000
32	2	10	2000
48	3	15	3000
64	4	20 (full)	4000

PRESSURE SPRAYERS: Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated.

Amt. Lawn Weed Killer to Use:		Amt. of Water in Sprayer (Gallons)		Area to be Sprayed: (Sq. Feet)
Tbsp.	Fl. Oz.			
6	3	3		200
12	6	1½		400
24	12	3		800



DM-2000 LAWN WEED KILLER

Contains Trimec® Herbicide⁽¹⁾

KEEP FROM FREEZING

1 QUART COVERS 2000 SQ. FT.

Contains: 2,4-D, MCPP and Dicamba

Also Kills: DANDELION, KNOTWEED, HENBIT, PLANTAINS, SPURGE, POISON IVY,
and most other broadleaf weeds.

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	1.31%
**Dimethylamine salt of 2,4-dichlorophenoxy acetic acid	2.88%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	0.30%
Dimethylamine salts of related compounds	0.04%

INERT INGREDIENTS	95.47%
TOTAL	100.00%

*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent to 0.091 pounds per gallon, equivalent to 1.079%.

**2,4-dichlorophenoxyacetic acid equivalent to 0.201 pounds per gallon, equivalent to 2.391%.

***3,6-dichloro-o-anisic acid equivalent 0.021 pounds per gallon, equivalent to 0.248%.

⁽¹⁾Trimec® is a Reg. T.M. of PBI-Gordon Corp., U.S. Patent No. 3,284,186.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See right panel for additional cautions

SOLD BY

L. B. GARDNER, INC.
1429 FAIRMOUNT AVE. N.W.
ATLANTA 16, GEORGIA

NOTE: We
sprayer, or
can drift
HOW TO
To spray
do not u
tbsp.) in
and spr
"wet" ec
WEEDS
may be
Zoysia
other si
such as
SPOTTED
and SH
Lawn W
leafed la
Plantain,
CONTROL
Bedstraw
Black Medic
Buckhorn
Chicory
Chickweed
Clover
Dandelion
Dock
Ground Ivy
Heal-All

CAUTION:
vegetab
shrubs, tr
not conta
Do not pe
Do not st
insecticid
Carpeter
where d
present. V
oughly a
lawn spr
are less li
underlaid
shrubs. T
Augustine

CAUTION:
skin, eye,
contact, f
tion of fe
NOTICE:
conforms
label. S
weather,
and cond
vary, sell
stated on
CONTROL
NUMBER

E.P.A. Reg

ACCEPTED
 Jul 20, 1972
 UNDER THE FEDERAL INSECTICIDE,
 FUNGICIDE AND WEED KILLER ACT
 FOR ECOLOGICALLY POISONOUS REGISTERED
 UNDER NO. 2155-61

DM-2000 LAWN WEED KILLER

Contains Trimec® Herbicide⁽¹⁾
KEEP FROM FREEZING
 1 QUART COVERS 2000 SQ. FT.
 Contains: 2,4-D, MCPP and Dicamba

Also Kills: DANDELION, KNOTWEED, HENBIT, PLANTAINS, SPURGE, POISON IVY,
 and most other broadleaf weeds.

ACTIVE INGREDIENTS:

- *Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 1.31%
- **Dimethylamine salt of 2,4-dichlorophenoxy acetic acid..... 2.88%
- ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)..... 0.30%
- Dimethylamine salts of related compounds..... 0.04%

INERT INGREDIENTS 95.47%
TOTAL 100.00%

^{*}2-(2-methyl-4-chlorophenoxy) propionic acid equivalent to 0.091 pounds per gallon, equivalent to 1.079%.
^{**}2,4-dichlorophenoxyacetic acid equivalent to 0.201 pounds per gallon, equivalent to 2.391%.
^{***}3,6-dichloro-o-anisic acid equivalent 0.021 pounds per gallon, equivalent to 0.248%.
⁽¹⁾Trimec is a Reg. T.M. of F&I-Gordon Corp., U.S. Patent No. 3,284,186.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See right panel for additional cautions

SOLD BY
 I. BENNETT, INC.
 1429 FARMGATE AVE. N.W.
 ATLANTA 18, GEORGIA



NOTE: Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.
HOW MUCH TO USE: SPOT SPRAY
 To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in 1/2 gal. water in pressure sprayer and spray as required, being certain to "wet" each weed.

WEEDS CONTROLLED: Lawn Weed Killer may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas with hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, Lawn Weed Killer controls most other broad-leaved lawn weeds, including Dandelion and Plantain, and a partial list is given below:

CONTROLS:		
Bedstraw	Henbit	Poison Ivy
Black Medic	Knotweed	Pursh
Buckhorn	Lambquarters	Ragweed
Chicory	Lespedeza	Shepherd's Purse
Chickweed	Mallow	Speedwell
Clover	Morning Glory	Spurge
Dandelion	Papergrass	Wild Carrot
Dock	Pigweed	Wild Lettuce
Ground Ivy	Plantain	Wild Onion
Heat-All	Poison Ivy	Yarrow

CAUTION: Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on Carpetgrass, dichondra or lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

CAUTION: Harmful if swallowed. Avoid skin, eye, and clothing contact. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.

CONTROL NUMBER 8209-572
 GAL. 8209-572
 E.P.A. Reg. No. 2155-61