

DIRECTIONS FOR USE: GENERAL INFORMATION: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This broadleaf weed killer is specifically formulated for maximum safety to seeding grasses and 2,4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill, the higher rates should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur. To insure that weeds are growing and able to absorb this product, irrigate the lawn 2-3 days prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70° F, sprays can be applied any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dichondra and most ornamental and vegetable crops. Amount of water in spray is not important, except to evenly distribute this product. If directions call for 16 oz. in 2 gallons per 1,000 square feet, it may be more convenient to use 16 oz. in 4 gallons depending on the sprayer. Generally, large volumes of water result in more uniform coverage.

WEED CONTROL: This product is especially effective against weeds such as Sheep Sorrel, Knotweed, Spotted Spurge, Chickweed, Oxalis and Henbit. In addition, it kills virtually all broadleaf weeds, some of which are listed below:

BEAN STRAW	GALINSOGA	PEPPERGRASS	SPURGE
BLANKET FLOWER	GROUND IVY	PIGWEEED	VELVET LEAF
BUCKHORN	HAWTHORN	PLANTAIN	WILD GARLIC
BURCHARD	HEAL ALL	POISON IVY	WILD GARLIC
CHICKWEED	HENBIT	POISON OAK	WILD RYE
CLOVER	KNOTWEED	RHUSLEAF	WILD RYE
CORN COCKLE	LAMBQUARTER	RAGWEED	WILD RYE
CRESS	LESPEDEZA	SHEEP SOREL	WILD RYE
CYPERUS	MALLOWS	SHEPHERD'S PURSE	WILD RYE
DANDELION	MORNING GLORY	SPEEDWELL	WILD RYE
DOCK	MUSTARD	TANSY MUSTARD	WILD RYE
	OXALIS	THISTLE	MORE

HOW TO MIX: 1. For use on St. Augustine grass, bentgrass and buffalo grass, mix 16 oz. in 2.4 gallons of water and uniformly spray 1,000 square feet. For example, an area 10 feet long and 100 feet wide. If necessary, repeat treatment in 2-3 weeks. 2. For use on most other grasses such as bermuda, bahia, bluegrass, centipede, fescue and rye, mix 20 oz. in 2.4 gallons of water and spray uniformly over 1,000 square feet. On hard to kill species, older weeds or under poor growing conditions, mix 32 oz. in 2.4 gallons of water and spray on 1,000 square feet with special attention to thorough coverage.

St. Augustine Grass Broadleaf Herbicide

Contains Trimec[®] Herbicide⁽¹⁾

Kills: Dandelions, Knotweed, Henbit, Chickweed, Plantains, Spurge and many other broadleaf weeds.

ONE QUART COVERS UP TO 2,000 SQUARE FEET
KEEP FROM FREEZING

ACTIVE INGREDIENTS

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	6.63
**Dimethylamine salt of 2,2-methyl-4-chlorophenoxypropionic acid	2.12
***Dimethylamine salt of D-cis-3,6-dichloro-2-naphthoic acid	0.25

INERT INGREDIENTS

12,4-dichlorophenoxyacetic acid equivalent 0.44 pounds per gallon or 0.52%	TOTAL	109.90
**2,2-methyl-4-chlorophenoxypropionic acid equivalent 0.145 pounds per gallon or 0.172%		
***3,6-dichloro-2-naphthoic acid equivalent 0.118 pounds per gallon or 0.143%		

Trimec[®] is a registered trademark of PBI-Gordon Corporation, U.S. Patent No. 3,284,186

FIRST! READ ENTIRE LABEL. OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY. USE IN CONFORMANCE WITH FEDERAL AND STATE REGULATION.

ACCEPTED
SEP 11 1987
EPA Reg No. 2155-62

KEEP OUT OF REACH OF CHILDREN.
CAUTION
SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS

FOR COMMERCIAL AND INDUSTRIAL USE ONLY, OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS

MANUFACTURED BY
NOT FOR SALE TO OR USE BY HOMEOWNERS
P. SCHNEITZ, INC.
140 CALHOUN AVE. NW
ATLANTA, GA 30303 404-521-1601

Delete
Addition

BEST AVAILABLE COPY

PRECAUTIONS FOR USAGE:

Only use lawn sprayers. Avoid drifts of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Coarse spray is less likely to drift. Do not apply to areas uncerlaid by roots of desirable trees and shrubs. Do not contaminate domestic or irrigation water.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.
IF SWALLOWED: Drink 1-2 glasses of water and induce vomiting by touching back of throat with finger. Never give anything by mouth to an unconscious person. Call a physician.
IF IN EYES: Flush with plenty of water. Call a physician.
IF ON SKIN: Wash with plenty of soap and water. Get medical attention.
IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply directly to water.

STORAGE AND DISPOSAL

STORAGE: Do not store with fertilizers, seeds, insecticides or fungicides.
PROHIBITION: Do not contaminate water, food or feed by storage or disposal or cleaning of equipment. Open dumping prohibited.
PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

EPA Reg No. 2155-62
EPA Est. No. 2155-GA-1

6673
101786

CONTROL NUMBER

NET CONTENTS GALLONS