DIRECTIONS

For CLEANING and DISINFECTING floors, walls. surfaces as found in hospitals, institutions, public rooms, industrial areas, and conveyances:

- 1. Apply by mop or cloth a solution of at least 3 ozs, in one gallon of water.
- In instances where surfaces are heavily contaminated with soil and/or organic matter, apply solution twice to insure disinfection.
- 3. If surfaces are very porous use a solution of at least 4 ozs, in one gallon of water.

For CLEANING and SANITIZING food handling equipment, apply by mop or cloth a solution of at least 2 oz. in one gallon of water. After sanitizing thoroughly, rinse with potable water before reuse.

For CLEANING and DEODORIZING apply by mop or cloth a solution of at least 1 oz. in one gallon of water.

NOTE: Product is a concentrate and must be diluted before use. Do not mix this product with soap or anionic detergents.

This product has been tested in accordance with the N. F. P. A, booklet 56, section 6-2 that outlines the testing procedure for testing the resistance of conductive floors. When used as directed, it will pass these requirements and can be used on conductive flooring in hospital operating rooms.

E.P. A. Reg. No. 2155-67 E.P.A. EST. No. 2155 - GA. 1 MOCK 2 DUAL-27 22, JULY 1982 MULTI-PURPOSE CLEANER, DISINFECTANT SANITIZER
AND DEODORANT

Phenol Coefficient: 27 vs. S. aureus (AOAC Tes: Method) 16.7 vs. S. typhosa

DANGER:

KEEP OUT OF REACH OF CHILDREN. See Right Panel for Additional Precautions.

FOR COMMERCIAL AND INDUSTRIAL USE ONLY, OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS.

Manufactured For

I. SCHNEID, INC.

: ARTICH STATION ATLANTA, GEORGIA 30318 floors, walls, tions, public

of at least 3

contaminat-

tion of at

od handling Jution of at er sanitizing ere reuse.

apply by in one gallon

must be di-

nce with the los the testing incrive floors, irrements and ital operating

2 DUAL-27

ACCEPTE DUAL - 27

SEP 20 1082 DUAL - 27

Logistored and Repartment of the Pesal And Purpose CLEANER OF THE MULT PURPOSE CLEANER, DISINFECTANT SANITIZER AND DEODORANT

Phenol Coefficient: 27 vs. S. aureus (AOAC Test Method) 16.7 vs. S. typhosa

ACTIVE INGREDIENTS

n Alkyt (60° C12, 30° C14, 5° C16, 5° C18) dimethyl benzyl ammonium chlorides

n Alkyt (60° C12, 32° C14) dimethyl ethylbenzyl ammonium chlorides

1.65°
Trisodium phosphate

1.65°
Tetrasodium ethylenediamine tetraacetate

1.00
Ethyl alcohol

INERT INGREDIENTS

Total Ingredients

1.000%

DANGER:

KEEP OUT OF REACH OF CHILDREN.
See Right Panel for Additional Precautions.

FOR COMMERCIAL AND INDUSTRIAL USE ONLY, OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS.

Manufactured For

I. SCHNEID, INC.

P. G. BOX 93180 AMERICH STATION ATLANTA, GEORGIA 30318 RECOMMENDED for cleaning, disinfecting and deodorizing in hospitals on hard surfaces such as floors, walls, ceilings, equipment and fixtures in areas such as operating rooms, clinics, isolation wards, nurseries, diet kitchens, and rest rooms.

RECOMMENDED for cleaning, disinfecting and deodorizing in schools and public institutions all hard surfaces such as corridors, classrooms, offices, rest rooms, locker rooms, showers; in fact, in all public places where cleanliness and bacterial control of environmental surfaces are vital.

RECOMMENDED for cleening, senitizing and deodorizing food handling equipment in food, meat, milk and beverage processing plants, restaurants and farms. Such equipment would include blanchers, evaporators, homogenizers, holding tanks, pipes, valves, utensils, etc. All equipment should be flushed with clear potable water prior to use.

In use dilutions, this product can be used on all hard inanimate surfaces not harmed by contact with water.

DANGER:

KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes eye damage. May cause skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Rinse empty container thoroughly with water and discard it.

CONTROL NUMBER CONT.