

ALGAECIDE 1250

Concentrated Swimming Pool Algaecide

KEEPS POOL WATER CLEAR AND FREE OF VISIBLE ALGAE

CONTROLS ALGAE SLIMES

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides	5.0%
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	5.0
Ethyl alcohol	2.5

INERT INGREDIENTS:	37.5%
	<u>100.0%</u>

DANGER: KEEP OUT OF REACH OF CHILDREN.
See Other Precautionary and First Aid Statements on side panels.

FOR INSTITUTIONAL OR INDUSTRIAL USE ONLY.

SOLD BY:
I. SCHNEID, INC.
P. O. BOX 93188
MARTECH STATION
ATLANTA, GEORGIA 30318

ACCEPTED

APR 25 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2155-73

This Algaecide may be used to sanitize pool areas such as walkways, diving boards, ladder rails, and walls and floors in shower and dressing areas. Prepare a sanitizing solution by mixing 1 ounce of this Algaecide with 4 gallons of water. Apply with mop, cloth, or sponge to those previously cleaned areas to be sanitized.

This Algaecide is effective against *Trichopython interdigitale*, a causative agent of athlete's foot, at a dilution of 1 3/4 ounces per gallon of water.

This Algaecide is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners, or premixed with other water treating chemicals. Algaecide works best when added by itself directly to pool water. This Algaecide at use is noncorrosive to inanimate surfaces.

DANGER

Keep Out of Reach of Children. Corrosive, causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.


Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

NOTE TO PHYSICIAN: Probable muscosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

CONTROL NUMBER

E.P.A. Reg. No. 2155-73 -
E.P.A. EST. 2155-GA. -1

CONT GAL.



KERR-McGEE

HERB-ALL*

POST EMERGENCE HERBICIDAL OIL

Jul - 11 - 1964 11342-927
2392-927

WEED CONTROL IN COTTON FIELDS

RECOMMENDED USAGE: When the cotton is 2"-2 1/2" high post emergence spray for weed control may be started at the rate of about 6-gallons per acre not more frequently than five-day intervals. A total of three such applications should be sufficient. Examine cotton plants carefully and if any evidence of scars or other damage is noted discontinue spraying *HERB-ALL until the damaged parts heal. Discontinue spraying when true bark begins to form or when small cracks begin to appear on the stem.

WEED CONTROL IN SOYBEAN FIELDS

RECOMMENDED USAGE: Application should be made between the twelfth and twenty-first days after emergence of the soybean plant. Use spray at the rate of five (5) gallons per acre. Only a single application is recommended, and it should be during this period. The same methods of application can be used for soybeans as for cotton.

WEED CONTROL IN CARROT PATCHES

RECOMMENDED USAGE: Carrots should not be sprayed until the plants have developed two fern-type leaves. After carrots reach the two fern leaf stage they may be sprayed as often as necessary, but not more frequently than once every seven days. Best results are obtained when HERB-ALL is applied at a rate of 30-40 gallons per acre. Do not spray carrots later than six weeks before harvest. HERB-ALL does not affect carrot growth or leave a residual odor or taste.

HERB-ALL is most effective when used in the early morning while the dew is still on the plants. Good results are obtained if the spray is applied on humid days at temperatures of 65-75 F; it is less effective under other conditions. Do not apply on windy days because the spray tends to drift and the application is spotty.

WEED CONTROL IN CRANBERRY BOGS

RECOMMENDED USAGE: HERB-ALL should be applied before the weeds are three inches tall. Sufficient spray must be used to thoroughly soak all weeds. This requires a minimum of 300 gallons

per acre and may require as much as 600 gallons per acre, depending upon the number of weeds present and density of vines. Best results are obtained when HERB-ALL is applied on a still, humid day when the temperature is below 75 F. Applications made during a light rain or on a foggy day, are particularly effective.

Make only one application per season. It may be applied while cranberry bushes are still dormant using an overhead spray, or HERB-ALL must be applied from a ground boom after new growth appears. HERB-ALL does not harm the woody parts of cranberry bushes but does have a detrimental effect on new growth.

The proper equipment, calibration, and a complete description of methods of application for weed control in crops listed on this label are shown in our booklet "Kerr-McGee HERB-ALL." To get this FREE booklet see your dealer or write direct to the address on this label.

Due to soil conditions, methods of application, weed conditions, and other factors beyond our control, no responsibility is assumed by Kerr-McGee for failure or damage to crops.

ACTIVE INGREDIENT: 100% PETROLEUM SOLVENT

Net weight 6.6 pounds per gallon. Bulk shipment for Herbicidal use.

FLAMMABILITY: Flash point over 100 F
RED LABEL NOT REQUIRED

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid prolonged or repeated contact with skin. Do not breathe vapors. May be flammable. Do not use or store near open flames. Use only when there is no hazard of drift. Harmful or fatal if swallowed; if swallowed do not induce vomiting and call physician immediately.



KERR-McGEE CORPORATION
Kerr-McGee Building
Oklahoma City, Oklahoma 73102

USDA Reg. No. 6737-4