## APPLICATION

This product is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime and other deposits. An initial slug application of 13.5 to 36.3 fluid ounces of concentrate per 1000 gallons of water to provide a concentration of 96 to 240 parts per million of this product based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 3.6 to 36.3 fluid ounces of con: centrate per 1000 gallons of water (24 to 240 parts per million of concentrate) should be employed every 2 to 5 days as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

## TOWERCIDE 5


CAUTION:

KEEP OUT OF THE REACH OF CHILDREN

• • • •

FOR INSTITUTIONAL OR INDUSTRIAL USE ON

SOLD BY:



Percent by weight Poly[oxyethylene (dimethyliminio)ethylene(dimethyliminio)-ethylene dichloride] 95.0% Weight per gallon . . . 

UTION:

KEEP OUT OF THE REACH OF CHILDREN

FOR INSTITUTIONAL OR INDUSTRIAL USE DNIX.

**SOLD BY:** 

NOTCIDE AND RODENTICIDE ACT NUMBER

LA RODICINIC POISON REGISTER

E. P. A. Reg. No. 2155-76

**CAUTION:** 

KEEP OUT OF THE REACH OF CHILDREN.

Harmful if swallowed. Avoid contact with skin and eyes. In case of contact flush with plenty of water for at least 15 minutes. If eye irritation persists get medical attention. Avoid contamination of food.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment or disposal of waste. ED Use only as directed on label. Do not reuse container. Destroy when empty.

CONT.

GAL.