

**DIRECTIONS FOR USE**

**Initial Dose:** 1 gallon of concentrate to 25,000 gallons of water.

**Maintenance Dose:** 1 quart of concentrate to 25,000 gallons of water.

Initial dose is used when filling pool at beginning of swimming season. If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat this treatment procedure as necessary. When pool is free of visible algae, use the recommended maintenance dose.

Maintenance dose should be added at 3-5 day intervals. If high temperatures prevail or pool has unusually heavy use, add maintenance dose more frequently.

If make-up water is added at the rate of more than 10% per week of pool capacity, add 1 quart of concentrate for each 5,000 gallons of fresh make-up water added per week.

When swimming season is over add 1 gallon of concentrate to 25,000 gallons of water in treating water left in pool. This dose helps provide a measure of control of algae growth during the winter months. This treatment will also help the servicing of the pool before it is returned to use the following season.

**FISH AND/OR WILDLIFE PRECAUTIONS:** This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public waters. Apply this product only as specified on this label.

When this concentrate is used as directed, it will kill and prevent development of *Chlorella Pyrenoidosa* (green), *Phormidium inundatum* (black algae), *Phormidium retzii* (Square D algae), *Lyngbya versicolor* (bluegreen) and other swimming pool algae.

E.P.A. Reg. No. 2155-83-  
E.P.A. EST. No. 2155-GA-1

# ALGAEICIDE 625

## Concentrated Swimming Pool Algaecide

KEEPS POOL WATER CLEAR AND FREE OF VISIBLE ALGAE

CONTROLS ALGAE SLIMES

**ACTIVE INGREDIENTS:**

n-Alkyl(60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides	2.50%
n-Alkyl(68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	2.50%

<b>INERT INGREDIENTS:</b>	95.00%
<b>TOTAL</b>	<b>100.00%</b>

**WARNING: KEEP OUT OF THE REACH OF CHILDREN.**  
See Other Warnings and First Aid Statements on side panels.

**ACCEPTED**  
FOR INSTITUTIONAL OR INDUSTRIAL USE ONLY.

NOV 11 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2155-83

**SOLD BY:**  
**I. SCHNEID, INC.**  
P. O. BOX 93188  
MARTECH STATION  
ATLANTA, GEORGIA 30318

This concentrate may be used to sanitize pool areas such as walkways, diving boards, ladder rails, and walls and floor in shower and dressing areas. Prepare a sanitizing solution by mixing 1 ounce of the concentrate with 2 gallons of water. Apply with mop, cloth, or sponge to those previously cleaned areas to be sanitized, and wet thoroughly with the recommended dilution.

This concentrate is effective against *Trichopython interdigitale*, a causative agent of athlete's foot, at a dilution of 1 pint per 5 gallons of water (3.2 fluid ounces/gallon water).

This concentrate is compatible with most commonly used swimming pool chemicals, but should not be mixed with soap or other cleaners, or premixed with other water treating chemicals. This algaecide works best when added by itself directly to pool water. This concentrate at use is noncorrosive to inanimate surfaces.

**WARNING:**

Keep out of Reach of Children. Corrosive, causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

**FIRST AID**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid Alcohol. Call a physician.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

For small containers, rinse empty container thoroughly with water before discarding. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

**CONTROL NUMBER**

**NET CONTENTS GALLONS**