

SLO-MOWSHUN CONCENTRATE

A LIQUID GROWTH RETARDANT THAT INHIBITS GROWTH OF GRASS,
TREES, SHRUBS, AND IVY.

ALSO CONTROLS WILD ONION, GARLIC, AND BIENNIAL TYPE WEEDS.

ACTIVE INGREDIENTS:

Potassium salt of 6-hydroxy-3-(2H)-pyridazinone 30.2%

INERT INGREDIENTS: 69.8%

TOTAL 100.0%

This product contains 2.25 lbs. of Maleic Hydrazide per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING

HARMFUL OR FATAL IF SWALLOWED STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED - Call a Physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED - Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN - Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES - Flush eyes with plenty of water. Call Physician immediately.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

FOR COMMERCIAL/INDUSTRIAL ONLY. NOT FOR USE ON AGRICULTURAL ESTABLISHMENTS.

Manufactured For:

I. SCHNEID, INC.
1429 FAIRMONT AVENUE NW
ATLANTA, GEORGIA 30318-4153

GENERAL INFORMATION:

INHIBITS GRASS GROWTH: This liquid growth retardant is absorbed into the green grass blades and translocated by the plant fluids to the active growth areas of the plant where cell division is retarded. It also prevents seed head formation and slows down leaf growth. If an area is not mowed after the growth retardant is applied, the turf will slowly grow to 6 to 12 inches in height by the end of the growing season as the effect of the retardant gradually "wears off." This is especially effective where safety is a factor as it reduces the need for frequent mowing of hazardous locations and eliminates mowing of cuts and fills where a seasonal growth of 10 to 6-11 inches can be tolerated.

DIRECTIONS: Apply the Liquid Growth Retardant to green grass, since brown grass will not absorb the spray. One application per year made either early in the fall or in the spring, followed by correct maintenance procedures will produce required turf growth control.

If the product is not applied before the seed head is formed, the seed head will expand in spite of the presence of the concentrate. If this occurs, at least

one week should be allowed for the retardant to circulate throughout the plant. The area should then be mowed to no less than 4 inches in height. Inhibition of subsequent growth will then be observed.

If a neatly clipped appearance is required, the retarding application should be supplemented by mowing. Do not apply during the summer (June 1 - Sept. 25). (The major reason for summer mowing is the presence of annual grasses and weeds which will have to be controlled by mowing or with herbicides since they are not present when this concentrate is applied to perennial grasses.)

EFFECT OF RAIN ON TREATMENT: This retardant should be applied during a period when rain is not expected for 12 hours after treatment. Since this retardant is water soluble, a rain will wash the chemical off the leaf blade before absorption occurs. Once the retardant is absorbed, rain has no effect on performance.

This concentrate should be used on good turf, 3 or more years old, which needs frequent mowing, and areas that are difficult to mow.

CONDITIONS AFFECTING ABSORPTION OF CONCENTRATE BY GRASS: It has been found that a greater amount of concentrate will enter the leaf in the shortest time and give the best results when the plants are growing rapidly in moist soil. The concentrate is not absorbed well when 1) leaves are wilted and 2) during periods of extended growth when the soil is dry.

USE OF 2,4-D: 2,4-D may be used with this product when control of broadleaf weeds is desired in turf. A low volatile ester or amine salt formulation is compatible with this concentrate. In some areas weeds may emerge after the concentrate is applied. A second application of 2,4-D would be necessary to control these weeds. Observe all cautions and limitations on labeling of all products used in mixtures.

NOTES: 1) Do not pasture livestock in, or cut hay from, treated areas. 2) This product can be used along highways, airports, military installations, roughs of golf courses and similar areas. 3) At recommended application rates, this concentrate does not normally affect the color of grass. Under conditions of optimum absorption, color modification for a period of several weeks may be encountered. At times treated grass may be a greener color than untreated areas. It may also remain greener for a longer period under drought conditions. No adverse effects on color have been observed from fall application. Some delay in spring "green up" is normal. 4) Do not spray on trees when wind velocity exceeds 15 mph. 5) Use 2,4-D only on grasses recommended by the manufacturer's label. Follow manufacturer's label when handling or spraying 2,4-D to prevent drift.

STORAGE AND DISPOSAL

1.) Prohibition - DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE, DISPOSAL OR CLEANING OF EQUIPMENT. OPEN DUMPING PROHIBITED.

2.) Pesticide Disposal - Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

3.) Container Disposal - Plastic containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Metal containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS). WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Keep children and pets off treated areas until material is washed into the soil and grass is dry.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residue remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop or weeds blooming in the treatment area. Applications should be timed to provide maximum possible intervals. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment or washwaters.

It is a vic
with its la
Not for u
for comm
For use
crops, tin
or for col
The follo
if top ten
spray the
surfaces
gallons o
to the po
trees and
on applic
100 galk
water.
WHEN T
the spray
leaves h
pruning a
CONTRC
will inhibit
pruned in
on the rel
as soon a
Privet, P
Cissus, H
The follo
Oak, Sw

FALL AP
in 30-50
treatment
is inhibite
than untr
becomes
there is a
absorptio
biennial
SPRING
concentr
The two
ideal for
Dandelion
will absor
formation
period, a
appear
mowing
interfere
TO CON
ALONG H
to 2 galk
fall when
lawns on
seedlings
grains su
E.P.A. RE
EST. NO
CONTRC
NUMBER

BEST COPY AVAILABLE

MOWSHUN CENTRATE

DANT THAT INHIBITS GROWTH OF GRASS,
3, SHRUBS, AND IVY.
ON, GARLIC, AND BIENNIAL TYPE WEEDS.

hydroxy -3- (2H) - pyridazinone	30.2%
.....	69.8%
TOTAL	100.0%

contains 2.25 lbs. of Maleic Hydrazide per gallon.

UT OF REACH OF CHILDREN

WARNING

UL OR FATAL IF SWALLOWED
NT OF PRACTICAL TREATMENT

on Control Center immediately. Drink one or two glasses of water and
throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting
as person.

apply respiration if indicated.

ing and wash affected areas with soap and water.

or. Call Physician immediately.

R ADDITIONAL PRECAUTIONARY STATEMENTS

ONLY. NOT FOR USE ON AGRICULTURAL ESTABLISHMENTS.

Manufactured For:

I. SCHNEID, INC.
FAIRMONT AVENUE NW
ATA, GEORGIA 30318-4153

sorbed into
the active
o prevents
o 12 inches
o retardant
a factor as
ations and
up to 6-11

ass, since
made either
procedures

seed head
rs, at least

one week should be allowed for the retardant to circulate throughout the
plant. The area should then be mowed to no less than 4 inches in height.
Inhibition of subsequent growth will then be observed.

If a neatly clipped appearance is required, the retarding application should be
supplemented by mowing. Do not apply during the summer (June 1 - Sept. 25).
(The major reason for summer mowing is the presence of annual grasses and
weeds which will have to be controlled by mowing or with herbicides since
they are not present when this concentrate is applied to perennial grasses.)

EFFECT OF RAIN ON TREATMENT: This retardant should be applied
during a period when rain is not expected for 12 hours after treatment. Since
this retardant is water soluble, a rain will wash the chemical off the leaf blade
before absorption occurs. Once the retardant is absorbed, rain has no effect
on performance.

This concentrate should be used on good turf, 3 or more years old, which
needs frequent mowing, and areas that are difficult to mow.

CONDITIONS AFFECTING ABSORPTION OF CONCENTRATE BY GRASS: It has been found that a greater amount of concentrate will enter
the leaf in the shorter time and give the best results when the plants are
growing rapidly in moist soil. The concentrate is not absorbed well when 1)
leaves are wilted and 2) during periods of extended growth when the soil is
dry.

USE OF 2,4-D: 2,4-D may be used with this product when control of
broadleaf weeds is desired in turf. A low volatile ester or amine salt
formulation is compatible with this concentrate. In some areas weeds may
emerge after the concentrate is applied. A second application of 2,4-D would
be necessary to control these weeds. Observe all cautions and limitations
on labeling of all products used in mixtures.

NOTES: 1) Do not pasture livestock in, or cut hay from, treated areas. 2) This
product can be used along highways, airports, military installations, roughs
of golf courses and similar areas. 3) At recommended application rates, this
concentrate does not normally affect the color of grass. Under conditions of
optimum absorption, color modification for a period of several weeks may be
encountered. At times treated grass may be a greener color than untreated
areas. It may also remain greener for a longer period under drought
conditions. No adverse effects on color have been observed from fall
application. Some delay in spring "green up" is normal. 4) Do not spray on
trees when wind velocity exceeds 15 mph. 5) Use 2,4-D only on grasses
recommended by the manufacturer's label. Follow manufacturer's label
when handling or spraying 2,4-D to prevent drift.

STORAGE AND DISPOSAL

1.) **Prohibition - DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE, DISPOSAL OR CLEANING OF EQUIPMENT. OPEN DUMPING PROHIBITED.**

2.) **Pesticide Disposal -** Pesticide wastes are toxic. Improper disposal of
excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If
these wastes cannot be disposed of by use according to label instructions,
contact your State Pesticide or Environmental Control Agency, or the
Hazardous Waste representative at the nearest EPA Regional Office for
guidance.

3.) **Container Disposal -** Plastic containers: Triple rinse (or equivalent).
Then offer for recycling or reconditioning, or puncture and dispose of in a
sanitary landfill, or incineration, or, if allowed by state and local authorities,
by burning. If burned, stay out of smoke. Metal containers: Triple rinse (or
equivalent). Then offer for recycling or reconditioning, or puncture and
dispose of in a sanitary landfill, or by other procedures approved by state and
local authorities.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS). WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe
spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after
handling. Avoid contamination of food and feed. Keep out of reach of
domestic animals. Do not use on humans, household pets, or livestock.
Keep children and pets off treated areas until material is washed into the soil
and grass is dry.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and other wildlife. Birds feeding on treated
areas may be killed. Do not contaminate water by cleaning of equipment or
disposal of wastes. Do not apply when weather conditions favor drift from
treated areas. Apply this product only as specified on this label. This
pesticide is highly toxic to bees exposed to direct treatment or to residue
remaining on the treated area. Do not apply when bees are actively visiting
the crop, cover crop or weeds blooming in the treatment area. Applications
should be timed to provide maximum possible intervals. Do not apply directly
to water, to areas where surface water is present or to intertidal areas below
the mean high water mark. Do not contaminate water by disposal of
equipment washwaters.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent
with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or
for commercial seed production, or for research purposes.

For use on plants in non-crop and non-lumber areas only. Not for use on
crops, timber, or other plants being grown for sale or other commercial use,
or for commercial seed production, or for research purposes.

The following spray solution should be sprayed on the upper half of the tree
if top terminal growth is to be inhibited. If the side of a tree is to be inhibited
spray the half of the tree on the pruned side. For shrubs and ivy, spray all
surfaces of the leaves and bark. Add 1 1/3 gallons of concentrate to 100
gallons of water. Spray this concentrate to wet all surfaces of leaves and bark
to the point of drip (not run off). Hose on attachments can be used to spray
trees and shrubs. Wet all surfaces of leaves and bark. Consult instructions
on applicator for delivering the equivalent of 1 1/3 gallons of concentrate in
100 gallons of water (equals 4 tablespoons) of concentrate in 1 gallon of
water.

WHEN TO SPRAY TREES AND SHRUBS: Best results are obtained when
the spray is applied to vigorous new growth in 1) Spring, just as the new
leaves have expanded and new growth has started; 2) Summer, after
pruning and new regrowth is 2-4 inches in length.

CONTROL OF TREES, SHRUBS, AND IVY: This liquid growth retardant
will inhibit growth of various trees, shrubs, and ivy. The plants should first be
pruned into the desirable shape. After regrowth of 2-4 inches occurs, spray
on the retarding concentrate. This concentrate can also be applied in spring
as soon as new leaves have expanded. The following shrubs can be treated:
Privet, Pyracantha, Myrtus, Xylomas, Viburnum, Eugeniz, Pittosporum
Cissus, Hahns Ivy, Algerian Ivy, Honeysuckle, Forsythia and Ice Plant.
The following tree species can be treated: Sycamore, Willow, Alder, Poplar,
Oak, Sweet Gum, Elm and Tamarack.

FALL APPLICATION: (Oct. 1 - Nov. 15) Use 1 1/3 gallons of the concentrate
in 30-50 gallons of water per acre to reduce growth next season. The fall
treatment is applied to green grass before it becomes dormant. Since growth
is inhibited the following spring the area will "green up" about two weeks later
than untreated grass. Apply the retardant as close to the time the grass
becomes dormant as possible but when it is still green. Do not spray when
there is a cover of fallen leaves as this will interfere with coverage and
absorption, causing poor results. This product will control wild onions and
biennial type weeds such as Dandelion and Plantain when applied in the fall.
SPRING APPLICATION: (About April 10 - June 1) Use 1 1/3 gallons of the
concentrate in 50 gallons of water per acre to reduce or eliminate mowing.
The two week period in the spring when the grass has grown 2-4 inches is
ideal for applying the concentrate. This is the period when Forsythia and
Dandelion are in full bloom. The grass is in a vigorous growing condition and
will absorb the chemical at the maximum rate. It will prevent seed head
formation and inhibit growth of foliage. If spraying is delayed beyond this
period, a mowing may be required after application to trim the area to improve
appearance. Allow one week for translocation of the chemical before this
mowing. Do not spray newly mowed areas covered with debris, this will
interfere with coverage and absorption, causing poor results.

**TO CONTROL WILD ONIONS AND WILD GARLIC IN GRASS AREAS
ALONG HIGHWAYS AND IN LAWNS** (not pasture or crop land): Apply 1
to 2 gallons of concentrate to 20 to 100 gallons of water per acre in spring or
fall when about 90-95 percent of the bulbs have started regrowth. Use on
lawns only when grass is brown and dormant. Do not apply to young
seedlings less than one season old or winter pastures consisting of small
grains such as rye.

E.F.A. REG. NO. 2155-105-
EST. NO. 2155-GA-1

6534/777777

CONTROL
NUMBER

NET CONTENTS
GALLONS

BEST COPY AVAILABLE

343

ACCEPTED
with COMMENTS
In EPA Letter Dated:

APR 11 1995

For the Federal Insecticide
Fungicide and Rodenticide Act
Registration No. 2155-105

2155-105

4
3
2
1

BEST COPY AVAILABLE

3