

PM 31 2155-117 10/2
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

I. Shneid, Inc.
1429 Fairmount Ave. NW.
Atlanta, GA 30381-46501

JUN 18

Attention: Gerard Longtin

Gentlemen:

Subject: BR-62B
EPA Registration No. 2155-117
Your Submission Dated March 15, 1991
EPA Received Date March 19, 1991

The amendment referred to above, submitted in connection with registration under FIFRA, is acceptable provided that you submit five (5) copies of your final printed labeling before you release the product for shipment.

The submitted data are acceptable to support the formula change. The revised Confidential Statement of Formula with changes in the inert ingredient has been reviewed and found to be acceptable. The amendment request to add additional organisms to your label is acceptable.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

If you have any questions concerning this letter, please contact Karen M. Leavy at (703)-557-3966.

Sincerely yours,



John H. Lee
Product Manager (31)
Antimicrobial Program Branch
Registration Division (H7505-C)

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

BEST AVAILABLE COPY

BR-62B

ONE STEP GERMICIDAL DETERGENT AND DEODORANT


Pseudomonacidal • Staphylocidal • Salmonellacidal
• Fungicidal • Virucidal

ACTIVE INGREDIENTS:	
Didecyl dimethyl ammonium chloride	4.61%
n-Alkyl (C14 50%, C12 40%, C10 10%)	
dimethyl benzyl ammonium chloride	3.07%
INERT INGREDIENTS:	92.32%

KEEP OUT OF REACH OF CHILDREN
DANGER

Manufactured for:

I. SCHNEID INC.
1429 FAIRMONT AVE., N.W.
ATLANTA GA 30381-4601



A concentrated, multi-purpose germicidal detergent and deodorant effective in hard waters up to 400 ppm hard water (calculated as CaCO₃) plus 5% organic serum. Disinfects, cleans, and deodorizes in one labor saving step. Recommended for use in hospitals, nursing homes, schools, colleges, veterinary clinics and animal life science laboratories and federally inspected meat and poultry establishments, equine farms, tack shops, pet shops, airports, kennels, hotels and motels, poultry and turkey farms, dairy farms, and hog farms, breeding and grooming establishments.

Bactericidal @ 1:128 dilution (1 ounce per gallon of water) against the following organisms according to the current AOAC. Use dilution test method, modified in the presence of 400 ppm synthetic hard water (calculated as CaCO₃) plus 5% organic serum.

<i>Pseudomonas aeruginosa</i>	<i>Mycobacterium</i>	<i>Enterobacteriaceae</i>
<i>Staphylococcus aureus</i>	<i>Proteus</i>	<i>E. coli</i>
<i>Salmonella</i>	<i>Shigella</i>	<i>Streptococcus aureus</i> (antibiotic resistant)
<i>Enterobacteriaceae</i>	<i>Shigella flexneri</i>	<i>Streptococcus faecalis</i> (antibiotic resistant)
<i>Streptococcus pyogenes</i>	<i>Shigella sonnei</i>	<i>E. coli</i> (antibiotic resistant)
<i>Streptococcus lactis</i>	<i>Shigella dysenteriae</i>	<i>Staphylococcus epidermidis</i> (antibiotic resistant)
<i>Enterobacteriaceae</i>	<i>Shigella flexneri</i>	<i>Staphylococcus saprophyticus</i> (antibiotic resistant)
<i>Salmonella typhi</i>	<i>Shigella flexneri</i>	<i>Pseudomonas aeruginosa</i> (antibiotic resistant)
<i>Salmonella typhimurium</i>	<i>Shigella flexneri</i>	

Fungicidal against *Trichophyton menthagale* and *Candida albicans* according to the AOAC Fungicidal Test, modified in the presence of 400 ppm hard water (calculated as CaCO₃) plus 5% organic serum @ 1:128 dilution.

Virucidal against the following viruses according to the virucidal qualification, modified in the presence of 400 ppm hard water plus 5% organic serum @ 1:128 dilution. (1 oz. per gal. of water).

<i>Coxsackie B virus</i>	<i>Poliovirus</i>	<i>Adenovirus type 4</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 1</i>	<i>Poliovirus type 2</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 2</i>	<i>Poliovirus type 3</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 3</i>	<i>Poliovirus type 4</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 4</i>	<i>Poliovirus type 5</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 5</i>	<i>Poliovirus type 6</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 6</i>	<i>Poliovirus type 7</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 7</i>	<i>Poliovirus type 8</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 8</i>	<i>Poliovirus type 9</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 9</i>	<i>Poliovirus type 10</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 10</i>	<i>Poliovirus type 11</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 11</i>	<i>Poliovirus type 12</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 12</i>	<i>Poliovirus type 13</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 13</i>	<i>Poliovirus type 14</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 14</i>	<i>Poliovirus type 15</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 15</i>	<i>Poliovirus type 16</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 16</i>	<i>Poliovirus type 17</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 17</i>	<i>Poliovirus type 18</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 18</i>	<i>Poliovirus type 19</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 19</i>	<i>Poliovirus type 20</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 20</i>	<i>Poliovirus type 21</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 21</i>	<i>Poliovirus type 22</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 22</i>	<i>Poliovirus type 23</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 23</i>	<i>Poliovirus type 24</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 24</i>	<i>Poliovirus type 25</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 25</i>	<i>Poliovirus type 26</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 26</i>	<i>Poliovirus type 27</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 27</i>	<i>Poliovirus type 28</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 28</i>	<i>Poliovirus type 29</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 29</i>	<i>Poliovirus type 30</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 30</i>	<i>Poliovirus type 31</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 31</i>	<i>Poliovirus type 32</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 32</i>	<i>Poliovirus type 33</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 33</i>	<i>Poliovirus type 34</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 34</i>	<i>Poliovirus type 35</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 35</i>	<i>Poliovirus type 36</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 36</i>	<i>Poliovirus type 37</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 37</i>	<i>Poliovirus type 38</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 38</i>	<i>Poliovirus type 39</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 39</i>	<i>Poliovirus type 40</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 40</i>	<i>Poliovirus type 41</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 41</i>	<i>Poliovirus type 42</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 42</i>	<i>Poliovirus type 43</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 43</i>	<i>Poliovirus type 44</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 44</i>	<i>Poliovirus type 45</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 45</i>	<i>Poliovirus type 46</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 46</i>	<i>Poliovirus type 47</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 47</i>	<i>Poliovirus type 48</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 48</i>	<i>Poliovirus type 49</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 49</i>	<i>Poliovirus type 50</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 50</i>	<i>Poliovirus type 51</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 51</i>	<i>Poliovirus type 52</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 52</i>	<i>Poliovirus type 53</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 53</i>	<i>Poliovirus type 54</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 54</i>	<i>Poliovirus type 55</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 55</i>	<i>Poliovirus type 56</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 56</i>	<i>Poliovirus type 57</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 57</i>	<i>Poliovirus type 58</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 58</i>	<i>Poliovirus type 59</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 59</i>	<i>Poliovirus type 60</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 60</i>	<i>Poliovirus type 61</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 61</i>	<i>Poliovirus type 62</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 62</i>	<i>Poliovirus type 63</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 63</i>	<i>Poliovirus type 64</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 64</i>	<i>Poliovirus type 65</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 65</i>	<i>Poliovirus type 66</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 66</i>	<i>Poliovirus type 67</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 67</i>	<i>Poliovirus type 68</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 68</i>	<i>Poliovirus type 69</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 69</i>	<i>Poliovirus type 70</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 70</i>	<i>Poliovirus type 71</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 71</i>	<i>Poliovirus type 72</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 72</i>	<i>Poliovirus type 73</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 73</i>	<i>Poliovirus type 74</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 74</i>	<i>Poliovirus type 75</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 75</i>	<i>Poliovirus type 76</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 76</i>	<i>Poliovirus type 77</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 77</i>	<i>Poliovirus type 78</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 78</i>	<i>Poliovirus type 79</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 79</i>	<i>Poliovirus type 80</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 80</i>	<i>Poliovirus type 81</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 81</i>	<i>Poliovirus type 82</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 82</i>	<i>Poliovirus type 83</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 83</i>	<i>Poliovirus type 84</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 84</i>	<i>Poliovirus type 85</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 85</i>	<i>Poliovirus type 86</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 86</i>	<i>Poliovirus type 87</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 87</i>	<i>Poliovirus type 88</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 88</i>	<i>Poliovirus type 89</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 89</i>	<i>Poliovirus type 90</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 90</i>	<i>Poliovirus type 91</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 91</i>	<i>Poliovirus type 92</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 92</i>	<i>Poliovirus type 93</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 93</i>	<i>Poliovirus type 94</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 94</i>	<i>Poliovirus type 95</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 95</i>	<i>Poliovirus type 96</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 96</i>	<i>Poliovirus type 97</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 97</i>	<i>Poliovirus type 98</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 98</i>	<i>Poliovirus type 99</i>
<i>Corona Parvovirus</i>	<i>Poliovirus type 99</i>	<i>Poliovirus type 100</i>

*Kills HIV-1 (AIDS virus) on pre-cleaned environmental surface objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus type 1 (HIV-1) (ASSOCIATED WITH AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gown, face mask, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures. Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of disinfectant.

Contact Time: Effective against HIV-1 (AIDS virus) on hard non-porous surfaces in the presence of a moderate amount of organic soil (5% blood serum) when used at a 1:128 dilution (providing 600 ppm of active quatamers) on 400 ppm CaCO₃ hard water for a contact time of 4 minutes at room temperature (20-25° C). Use a ten (10) minute contact time for disinfection against all other bacteria, fungi, and viruses claimed (listed on labeling).

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

EPA Reg No. 2155-117
EPA Est No. 2155-01-1

CONTROL
NUMBER

578920990
NET CONTENTS
GALLONS

DIRECTIONS
It is a violation of Federal Law to use this product.
GENERAL: USE DIRECTIONS FOR DISINFECTANT
For use on hard non-porous surfaces such as tile, grout, surfaces. Remove gross dirt and heavy soil deposit 1 ounce per gallon of water for a minimum contact time applied with a mop, sponge, or cloth as well as solution is prepared fresh for each use then discard to be washed or polished.
For use in meat and poultry plants as a general clean or mechanical cleaning device in all departments. If packaging materials must be returned from the compound, all surfaces in the area must be thorough of water.

STATEMENT OF PRACTICE
In case of contact, immediately flush eyes or skin with water. If swallowed, drink promptly a large amount of water. If not available, drink large quantities of water. A NOTE TO PHYSICIAN: Probable mucousal damage.

HAZARDS TO HUMANS AND ANIMALS
CORROSIVE: Causes eye damage and severe skin irritation. To protect eyes wear goggles or face shield handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.
STORAGE AND DISPOSAL

PROHIBITION
Do not contaminate water, food or feed by storage or disposal. Empty container.

PESTICIDE D. DISPOSAL: Pesticide wastes are acutely toxic. Do not mix with other pesticides, solvents, or food. Do not use according to label instructions, contact your local health department for disposal of these wastes. Agency, or the Hazardous Waste representative at your facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of in a sanitary landfill, or by incineration, or by other means approved by local, state, or federal authorities. Do not reuse or refill this container.

FIBER DRUMS WITH LINERS: Completely empty to avoid slinging particles. Empty residue into a sanitary landfill or by incineration if allowed by State and cannot be reused, dispose of in the same manner as above.

METAL CONTAINERS: Triple rinse (or equivalent) and dispose of in a sanitary landfill, or by incineration, or by other means approved by local, state, or federal authorities.

GENERAL: Consult, federal, state, or local disposal authorities for limited open burning.