



PM 03 2155-129 3-4-98 5531810 160 18 10/3

	<p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <u> x </u> Registration <u> </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 2155-129	Date of Issuance: MAR 4 1998
		Term of Issuance: Conditional	
		Name of Pesticide Product: General Purpose Liquid Insect Killer	
Name and Address of Registrant (include ZIP Code): John J. Walker, Ph.D. I. Schneid 1429 Fairmont Avenue, N.W. Atlanta, GA 30318-4153			
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none">1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.2. Make the following label changes:<ol style="list-style-type: none">a. Revise the EPA Registration Number to read, "EPA Reg. No. 2155-129".			
Signature of Approving Official: 		Date: MAR 4 1998	

EPA Form 8570-6

Bedwards A: 2155/129. reg 1/15/98

b. In the description for Indoor Application, add the statement "Not for use in Federally inspected meat and poultry plants."

c. Direct the user to read entire label by inserting the following statement at the top of the rear or side panel in bold, all caps.

"READ ENTIRE LABEL BEFORE EACH USE."

d. In the Directions for Use on DOGS, change the sentence "Not recommended for use on puppies less than four weeks old." to "Do not use on puppies under twelve (12) weeks old." per PR Notice 96-6.

e. Add the following statements to the Directions for Use on DOGS:

"Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals."

"Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately."

f. Every product label should have a company or Animal Poison Control Center, or other telephone number that consumers may call for the purpose of receiving information

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager 13
Insecticide Branch
Registration Division (7505C)

Enclosure

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying surface or surface applications. Do not use on humans. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. **PESTICIDE DISPOSAL:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility. **CONTAINER DISPOSAL:** CONTAINERS ONE GALLON AND SMALLER: Do not reuse container. Wrap container in several layers of newspaper and discard in trash. CONTAINERS LARGER THAN ONE GALLON: MEET CONTAINER'S Triple-rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **PLASTIC CONTAINERS:** Triple-rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. **GLASS CONTAINERS:** Triple-rinse or equivalent, then dispose of in a sanitary landfill or by other approved state and local procedures.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

SHAKE WELL BEFORE USING.

INDOOR APPLICATION

Do not use in food/feed areas of food/feed handling establishments, restaurants, or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared food is served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Nonfood/feed areas are areas such as garages, rooms, lawns, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). Vacuum treated areas and ventilate before occupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended. Do not apply to classrooms while in use.

SURFACE SPRAYING: TO CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDS, CENTIPEDS, SCORPIONS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, CRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES, AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests, and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices, and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Tilt beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as needed.


TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

MANUFACTURED BY:

I. Schneid

I. SCHNEID • P.O. BOX 93188 • ATLANTA, GA 30377-0188

6350/BULK/PL NEW



*Not for use
in Federally
inspected meat
and poultry
plants.*

GENERAL PURPOSE Liquid Insect Killer

Ready-to-Use Stable Emulsion Insecticide

- For control of fleas, brown dog ticks, and lice on premises and on dogs. Controls fleas and ticks on dogs for 35 days
- Aids in control of ticks on dogs for an additional 2 weeks
- For control of ticks which may cause Lyme disease on premises
- For use outdoors, in homes and non-food areas of food/feed processing establishments, kennels, commercial and office buildings, warehouses, theaters, schools, motels, and hotels

ACTIVE INGREDIENT:

*Permethrin 0.50%

INERT INGREDIENTS: 99.50%

TOTAL 100.00%

*[3-PHENOXYPHENYL] METHYL (+/-) CY/TRANS 3-(2, 2-DICHLOROETHOXY) 2, 2-DIMETHYL-4-PROPANECARBOXYLATE, CY/TRANS RATIO: MIN. 35% (+/-) CY AND MAX. 65% (+/-) TRANS.
EPA Reg. No. 2155-RED EPA Est. No. 2155-GA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE RIGHT SIDE PANEL FOR ADDITIONAL PRECAUTIONS.

FOR COMMERCIAL/AGRICULTURAL USE ONLY.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from at Apply spray liberally to empty chests, closets, bureaus, and other storage areas may be treated lightly, keeping nozzle at least three feet from fabric to kill LICE AND LOUSE EGGS. Spray in an inconspicuous area to test for spray entire area to be treated. Spray from a distance of 8 to 10 inches furniture that cannot be either laundered or dry cleaned. Allow all treated **SPACE SPRAYING:** Close doors and windows and shut off ventilating s for rapid control of HOUSE FLIES, FRUIT FLIES, GNATS, MOSQUITOES, SI **MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, A** uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic **FOODS:** To kill exposed or accessible stages of the above named craw Contact insects directly when possible. Keep area closed for 2 hours. **ON LIVESTOCK:** To protect cattle (beef and dairy), goats, sheep, hogs, **AND GNATS:** Apply a light mist sufficient to wet the surface of the hair. **TO CONTROL STABLE FLIES, HORSE FLIES, AND DEER FLIES:** Apply at a Repeat treatment daily or at intervals necessary to give continued protec **TO CONTROL BLOODSUCKING LICE:** Apply to the infested areas of the every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls, and nests or cages thori **TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES:** Spray crevices of r **ON DOGS:** To control FLEAS, TICKS, AND LICE: Start spraying at the tail body is covered, including the legs and underbody. While spraying, fluff thoroughly, but do not saturate animal. Do not spray into eyes or face. A product. Repeat as necessary.

NOTE: Not recommended for use on puppies less than four weeks old. Do not

OUTDOOR APPLICATION

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE: control TICKS including the DEER TICK (Ixodes dammini), LONG STAR (Amblyomma americanum), AMERICAN DOG TICK (Dermacentor variabilis) and WESTER

* For each area listed below, apply until surface is wet (approximately 1

* Treat around windows and doors, porches, screens, eaves, patios, garag

* Treat a band of vegetation and leaf litter 6 to 10 feet wide around and

* To help control ticks in outdoor activity areas, treat a band of vegetati

where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters, WHITEFLIES, APHIDS, EXPOSED THIRPS, SPIDERS, RED MITES, and LEAF

and for spot treatments of incipient insect infestations to prevent sporadic

violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, da

crassula, grape vines, Kente palms, laurels, marigolds, rhododendrons, r

EFFECTIVE WHEN USED AS DIRECTED AGAINST WHITEFLIES, APHIDS, SPI

PAVEMENT ANTS, CLOVER MITES, AND ARMYWORMS: When spraying

Be cautious about wetting tender foliage, young plants, and new growth

directly whenever possible. Repeat treatment as infestations occur.

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat

exceed 5 m.p.h., but do not apply when wind speeds exceed 10 m.p.h.

of 2.5 pints per acre (0.7 ounce per 1000 square feet). Using truck-mount

100 HT traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipm

water be treated with this product. Not to be used within 100 feet (30 m

Buyer assumes all risks of use, storage, or handling of this product not in

**CONTROL
NUMBER**

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 4 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2155-129

9.24.97, 7:34 PM