FOR GARDEN SOIL: To control wireworms, white grubs, Japanese beetle grubs, root maggots on radishes, onions and cabbage work into top 2 or 3 inches of soil when preparing seed bed. One pound will treat 1000 square feet. Remains effective all season.

FOR FOLIAGE DUST: To control grasshoppers, thrips

and plant bugs on rosebuds, flowers, shrubs and other ornamentals dust directly on the plant and insect. Repeat as necessary to prevent damage.

FOR BULB TREATMENT: Before storing bulbs and for treatment just before planting to control thrips on Dahlias, Gladiolus, Hyacinths, Tulips, and most flowering bulbs, place a few bulbs in a strong paper bag, dust a tablespoonful of 10% Chlordane Dust on the bulbs, close opening and then shake vigorously.

10

CRABGRASS CONTROL: Apply 15 lbs. to each 1000 square feet of lawn or turf to suppress crabgrass seed-tings before germinating or sprouting occurs. Use as an early season preventative treatment—not effective for control of growing crabgrass plants. Timing of application is important and should be applied before any germinating occurs, which in many areas is during the period April 15th to May 1st. To be safe apply before April 15th. This treatment will not burn or injure established lawns. Keep children and pets off treated areas until this material has been completely watered into the soil and the grass has completely dried. FOR LAWNS: To control ants, chinch bugs and chiggers, dust lightly at the rate of 1 pound to 1000 square feet of area. Repeat as necessary. For white

arubs Japanese beetle arubs and sod webward?

10%CHLORDANE

DUST

A HANDY GARDEN - LAWN - SOIL INSECTIGIDE

Total100%

*Equivalent to 6% Octachloro-4,7-Methano/Tetrachydroindane and 4% related compounds.

DIRECTIONS

IN THE GARDEN: To control cutworms, earwigs, mole crickets, sowbugs, millipedes, and leaf cutting ants dust liberally on the ground around corn, petatoes and ornamentals. Remains effective at least 10 days. Repeat as necessary. Do not

R

Dust on the builds, close opening and men snake vigorously.

CRABGRASS CONTROL: Apply 15 lbs. to each 1000 square feet of lawn or turf to suppress crabgrass seedtings before germinating or sprouting occurs. Use as an early season preventative treatment—not effective for control of growing crabgrass plants. Timing of application is important and should be applied before any germinating occurs, which in many areas is during the period April 15th to May 1st. To be safe apply before April 15th. This treatment will not burn or injure established lawns. Keep children and pets off treated areas until this material has been completely watered & into the soil and the grass has completely dried. FOR LAWNS: To control ants, chinch bugs and chiggers, dust lightly at the rate of 1 pound to 1000 square feet of area. Repeat as necessary. For white grubs, Japanese beetle grubs and sod webwork ? (lawn moth) apply liberally at the rate of one pound to 500 square feet. Water well with hose to get material into nests and living places. Effective entire season.

CAUTION: This material if taken internally can cause toxic symptoms. Avoid inhalation of dust and avoid contamination of foodstuffs or cooking utensils.

NON-WARRANTY: Neither the manufacturer nor the seller makes any representation or warranty, express or implied, and shall not be held responsible in any manner for any personal injury, property damage, or other type of loss resulting from the handling, storage and use of this product whether or not used in accordance with directions.

*Technical Chlordane 10%	7
Inert Ingredients 90%	
Total100%	influen.
*Equivalent to 6% Octachloro-4,7-Meth-	

*Equivalent to 6% Octachloro-4,7-Methano Tetrachydroindane and 4% related compounds.

£ --

DIRECTIONS

IN THE GARDEN: To control cutworms, earwigs, mole crickets, sowbugs, millipedes, and leaf cutting ants, dust liberally on the ground around corn, petatoes and ornamentals. Remains effective at least 10 days. Repeat as necessary. Do not feed treated forage to dairy animals or animals being finished for slaughter.

CAUTION:

Keep Out of Reach of Children. See other cautions on back panel.

USDA Reg. No. 2169-25

NET CONTENTS 8 OZ.

moderately beautiful represent affine

ACCEFTED