

DIRECTIONS FOR USE

GENERAL INFORMATION:

St. Augustine Broadleaf Weed Killer is specifically formulated for maximum safety to seedling grasses and 2, 4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill, the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions or in cases of accidental overdoses, some slight temporary discoloration of the St. Augustine foliage might occur.

To insure weeds are growing and able to absorb the herbicide, irrigate the lawn 2 or 3 days prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70 degrees F, sprays can be applied at any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dichondra and most ornamental and vegetable crops.

Amount of water in spray is not important except to evenly distribute the chemical. If directions call for 16 oz. in 2 gallons on 1000 square feet, it may be more convenient to use 16 oz. in 4 gallons on 1000 square feet, depending on the sprayer. Generally large volumes of water result in more uniform coverage.

HOW TO MIX:

- For use on St. Augustine grass, bentgrass and buffalo grass, mix 16 oz. in 2 to 4 gallons of water and uniformly spray 1000 square feet, for example, an area 10 feet wide and 100 feet long. If necessary, repeat treatment in 2 to 3 weeks, if hard to kill weeds persist.
- For use on most other grasses such as bermuda, bahia, bluegrass, centipede, fescue and rye, mix 20 oz. in 2 to 4 gallons of water and spray uniformly over 1000 square feet. On hard to kill species, or under poor growing conditions, or on older weeds, mix 32 oz. in 2 to 4 gallons of water and spray on 1000 square feet with special attention to thorough coverage of the hard to kill weeds.



St. Augustine Grass Broadleaf Herbicide

Contains Trimec® Herbicide⁽¹⁾

KILLS — DANDELIONS—KNOTWEED—HENBIT—CHICKWEED—PLAINTAINS—SPURGE—
and many other broadleaf weeds.

ONE QUART COVERS UP TO 2,000 SQUARE FEET
KEEP FROM FREEZING

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2, 4-dichlorophenoxyacetic acid	1.63
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	2.98
***Dimethylamine salt of Dicamba (3,6-dichloro-p-anisic acid)	2.25

INERT INGREDIENTS

2, 4-dichlorophenoxyacetic acid equivalent 0.044 pounds per gallon or 1.528%	21.94
2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 0.145 pounds per gallon or 1.72%	
3,6-dichloro-p-anisic acid equivalent 0.018 pounds per gallon or 0.21%	
TOTAL	100.0%

(1) Trimec® is a registered trademark of PBI Gordon Corporation, U.S. Patent No. 3,184,186

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONS

Sold by

QUANTICO, VIRGINIA

WEEDS CONTROLLED: St. Augustine Grass Herbicide is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Bedstraw	Galinsoga	Peppergrass	Spurge
Black Medic	Ground Ivy	Pigweed	Velvet Leaf
Buckhorn	Hawkweed	Plaintains	Wild Carrot
Bull Thistle	Heal-all	Poison Ivy	Wild Garlic
Burdock	Henbit	Poison Oak	Wild Lettuce
Buttercup	Knotweed	Purslane	Wild Onion
Chicory	Lambsquarter	Ragweed	Yarrow
Chickweed	Lespedeza	Sheep Sorrel	
Clover	Mallow	Shepherd's Purse	
Cocklebur	Morning Glory	Speedwell	And many
Dandelion	Mustard	Tansy Mustard	other broad
Fluck	Oxalis	Thistle	leaf weeds

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. If contact with eyes occurs flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not store in contact with fertilizers, seeds, insecticides or fungicides. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, seller makes no claims other than those stated on this label.

CONTROL
NUMBER

CONT.
GALS.

E. P. A. Reg. No. 211-102

PATTERSON'S®

SODIUM ARSENITE SOLUTION

**HERBICIDE, TREE KILLER, GRASSHOPPER
POISON AND TERMITE CONTROL**

ACTIVE INGREDIENT:

Sodium Arsenite (Na As O ₂)	40.0%
Total Arsenic expressed as metallic	23.1%
Arsenic in water soluble form expressed as metallic	23.06%

INERT INGREDIENTS 60.0%

DANGER: Keep out of reach of children.

Do not use nor store in or around the home.

See other warnings on side panel.

ANTIDOTE: Induce vomiting. Treat with mustard, magnesium or cup
ful with olive oil. If unconscious, give artificial respiration. CALL
A DOCTOR.

NET CONTENTS 1 GALLON