

### WOOD PRESERVATION

Pentachlorophenol Preservative can be effectively used to treat both old and new structures. When the wood is not in contact with the ground, as in the case of frames, sub-floors, sills, joists, etc., two applications of Pentachlorophenol Preservative should be made by swabbing or brushing. If individual members are being treated as in the case of a new building, it is best to soak the wood in Pentachlorophenol Preservative for from three to thirty minutes. All joints and ends of boards should be given thorough treatment.

Best results indicate that approximately one gallon of solution will be absorbed by 8 cubic ft. of wood. It is estimated one gallon will treat about 40 window sash 26"x26" size. When brushing or spraying, 2 gallons should be used to provide adequate protection.

Treated wood should be allowed to dry several days before being painted. Do not nail wall board or other absorbent material over the treated wood unless three or four days' drying period has elapsed.

Fence posts should be barked, seasoned and soaked for at least 24 hours and preferably 48 hours in a vertical position with butts downward. The best treatment is a full length coverage. If a butt treatment only is used, the posts should be treated to a length of three feet when the post is set two feet in the ground. Pines, red oak, basswood, black gum, and ash take treatment well; aspen and cottonwood do not treat well.

### TERMITE CONTROL

Spray or soak the soil, which has been removed from around the foundation of the building, to a depth of 12 inches, using 2 gallons of Pentachlorophenol Preservative per 5 cubic feet of soil or 2 gallons per 5 lineal feet of trench, or 5 square feet of ground surface. Direct the stream of liquid against the building below the

# PENTACHLOROPHENOL PRESERVATIVE

READY TO USE

AN EFFECTIVE  
WOOD PRESERVATIVE  
AND SOIL POISON

Also Controls Chicken Mites

### INGREDIENTS

ACTIVE	
Pentachlorophenol	5.00%
Petroleum Hydrocarbons	94.80%
INERT	0.20%
Total	100.00%

As defined by Federal Spec. TT-W-570.

### WARNING:

Keep out of reach of children.  
See other warnings on back panel.

NET CONTENTS      GALLON

### TERMITE CONTROL (Cont'd)

soil line. Replace the treated dirt, respraying at intervals as the earth is replaced. Decayed or termite attacked wooden sills should be replaced with new ones. Pentachlorophenol Preservative treated. Do not spray on plants.

### CHICKEN MITE CONTROL

The poultry house should be thoroughly cleaned including nests, dropping boards, roosts, floors, etc. When all parts are well dried, soak with Pentachlorophenol Preservative, using a spray or swab. Walls should be treated to a point somewhat higher than the roosts, or if the poultry house is heavily infested, entire walls and roofs should be included in the treatment. Be sure the solution is driven well into all cracks and joints.

After treatment, ventilate and dry for at least 48 hours before admitting chickens to the poultry house. Do not spray on chickens or other animals.

### NON-WARRANTY

Seller makes no warranty of any kind, either expressed or implied, concerning results to be obtained by the use of this product. Buyer assumes all risk of use or handling, whether or not in accordance with suggestions, recommendations, or directions of seller.

### CAUTION

Avoid excessive inhalation of vapors and contact with the skin. Keep away from foodstuffs, children, pets and live stock. In case of accidental swallowing, induce vomiting by administering emetic call a physician. To remove from skin, wash with plenty of soap and warm water.

### INFLAMMABLE

Contains volatile inflammable solvent. Do not pour, spill or spray near open flame and avoid breathing vapors.