

## DIRECTIONS

Continued

**Peaches:** To control early <sup>blight</sup> and late Blights, use 4 to 6 tablespoonfuls Bordeaux Mixture to 1 gallon water or 4 to 6 lbs. to 50 gal. water. Spray every 10 to 14 days after plants come up until just before harvest.

**Celery and Tomatoes:** To control early and late Blight, Leaf Spot use 4 to 5 tablespoonfuls to 1 gal. water or 4 to 5 lbs. to 50 gal. water, spraying or dusting plants in plant bed and at 10 day intervals in the field.

**Flowers:** To control Leaf Spot use 3 tablespoonfuls to 1 gallon of water or 3 lbs. to 50 gal. water and apply early in the season and repeat at 10 to 14 days intervals prior to blooming.

**Vines and Shrubs:** To control Leaf Spot use 2 tablespoonfuls to 1 gal. water or 2 lbs. to 50 gal. water by applying spray 3 times at weekly intervals when new growth appears in the spring.

**NOTICE:** Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

E.P.A. Reg. No. 2169-47 11-72

# BORDEAUX MIXTURE

### Active Ingredients:

Copper, expressed as metallic  
(in the form of Bordeaux  
Mixture) ..... 12.50%

Inert Ingredients ..... 87.50%

Total ..... 100.00%

### CAUTION:

#### Keep Out of Reach of Children.

Harmful if taken internally, drink large quantities of water, preferably with one ounce of Milk of Magnesia. Call a doctor immediately. Avoid breathing of dust.

Food utensils such as teaspoons or table-spoons should not be used for food purposes after use with pesticides. Do not reuse container. Destroy when empty.

**NET WEIGHT ONE POUND**

## DIRECTIONS

**Apple and Pear:** Delayed Dormant to control Scab, use 4 tablespoonfuls to 1 gallon or 4 pounds Bordeaux to 50 gallons oil spray.

**Pre-Blossom Spray.** One or two sprays to control Scab. Use 4 tablespoonfuls to one gallon or 4 pounds to 50 gallons of water.

**Late Summer Sprays** to control Scab, Blotch, Bitter Rot, Frog Eye, use 4 pounds Bordeaux Mixture to 50 gallons of water or with Summer Oil. It may be advisable to add 4 pounds of Hydrated Lime.

**Peaches and Plums:** A Dormant Spray to control Leaf Curl, San Jose Scale, Red Mite, use 3 tablespoonfuls to 1 gallon or 3 pounds to 50 gallon oil spray.

**Melons:** To control Anthracnose, Scab, Leaf Spot, use 3 tablespoonfuls to 1 gallon or 3 lbs. to 50 gal. of water. Spray as soon as plants appear and repeat at 5 day intervals during vining.

**Grapes:** To control Downy Mildew, Black Rot, use 3 to 4 tablespoonfuls of Bordeaux Mixture to 1 gallon or 4 lbs. to 50 gal water. Spray when shoots are 8 to 10 inches long, before blossom buds open, again after blossoms fall and again 10 to 14 days later.

**Cherries:** To control Leaf Spot, use 3 tablespoonfuls of Bordeaux Mixture to 1 gallon of water or 3 lbs. to 50 gal. water. On Sweet Cherries it is advisable to add 4 lbs. Hydrated Lime. Spray as soon as the petals fall and again one week later, plus two additional preharvest applications at 10-14 day intervals and two applications after fruit is harvested at 14 day intervals.