


PATTERSON'S  
1% LINDANE DUST

CONTROLS

Aphids, Thrips and Certain  
Worms and Beetles  
Also Roaches, Water Bugs, Silverfish

ACTIVE INGREDIENT:

Gamma isomer of benzene hexachloride from Lindane	1.0%
INERT INGREDIENTS:	99.0%
	100.0%



CAUTION:  
Keep Out of Reach of Children.  
See other cautions on side panel.

CONTENTS 1 POUND

MANUFACTURED BY, OR TO THE SPECIFICATIONS OF  
PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE

USDA Reg. No. 2169-60

KANSAS CITY, MISSOURI

DIRECTIONS

CRAWLING INSECTS: Roaches, waterbugs and silverfish may be controlled by 1% Lindane Dust. Thoroughly dust all hiding places which may harbor these insects, including cracks and crevices in foundation walls, openings in floors, areas beneath cabinets, sinks, stoves and refrigerators and similar places. Repeat application as often as needed for control.

AGRICULTURAL CROPS

HOME GARDENS: For insect control as outlined below, dust plants thoroughly, observing such restrictions as are indicated for certain crops.

LAWNS: Apply thoroughly for control of ticks and chiggers. Do not allow dust to contaminate garden crops.

CELERY: Apply 40 lbs. per acre to control aphid. Do not apply after plants are half grown or begin to bunch.

ASPARAGUS: Thoroughly dust at seedling and post cutting stages with 35 to 50 pounds per acre to control asparagus beetles and thrips.

Cole Crops (Cabbage, Broccoli, Brussels Sprouts): Apply about 30 pounds per acre to control thrips, aphids, and certain worms. Do not apply after edible parts start to form.

Cucurbits (Cucumbers, Squash, Pumpkins): Use 15 to 22 pounds per acre to control cucumber beetles. Do not apply after bloom. Apply 35 to 50 pounds per acre to control aphids, melon and pickle worms, and squash bug when insects first appear. Do not dust during harvest of pickles. Do not apply to squash after blossoms open.

Egg Plant: To control aphids, leaf miners, and lace bugs dust with

35 to 45 pounds per acre. Do not apply after blossoms fall.

Gladiolus: Apply 35 to 65 pounds per acre to control thrips.

Lettuce: Apply 30 to 40 pounds per acre to control aphids and certain worms. Do not apply to leaf lettuce after the seedling stage or to head lettuce after the heads begin to form.

Onions: Apply 35 to 65 pounds per acre to control thrips. Do not apply to spring or green onions.

Apples, Peaches, Plums, and Prunes: Use 35 to 65 pounds per acre to control aphids. Do not apply within 60 days of harvest.

Pears: To control pear psylla apply 35 to 65 pounds per acre. Do not apply within 60 days of harvest.

Peppers: Apply about 30 pounds per acre to control aphids, leaf miners, and foliage feeding worms. Do not apply after blossoms fall.

Spinach (seedling): To control aphids, leaf miners, thrips and web worms dust with 35 to 40 pounds per acre.

Strawberries: Apply about 15 pounds per acre to control spittle bugs and strawberry weevil. Do not apply after blossoms form.

Tomatoes: For the control of Colorado potato beetle, flea beetle, thrips and aphids dust with about 30 pounds per acre. Do not treat after fruit sets.

CAUTION

Avoid prolonged breathing of dust. May be absorbed through skin. Avoid contamination of feed and foodstuffs. Avoid contact with skin and eyes. Wash thoroughly after handling. Harmful if swallowed. Keep out of reach of children. Misuse as to quantity, timing, or method of application can cause damage or injury to animals, persons, property, crops or cause residues in excess of official tolerances.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Apply only as specified on this label.

Do not reuse container. Dispose in a safe place.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

