

RESISTANT OR EASY TO KILL SPECIES OF BRUSH

- | | |
|------------------------|--------------------------|
| —Easy to Kill— | IVY (Poison) |
| ALDER, SPECKLED | JUNEBERRY |
| BLACKBERRY | MAPLE (Soft) (Sugar) |
| BUCKBUSH | OAK (Black Jack, Burr, |
| CHERRY (Wild, Pin) | Ash (White) |
| CURRENT (Prickly, | Post White) |
| Sticky, Stink) | ORANGE (Orange) |
| DEWBERRY | PERSIMMON |
| ELM (American, Winged) | PLUM (Sand) |
| GOOSEBERRY | SUMAC (Smooth, Staghorn) |
| GRAPE | TRUMPET VINE |
| HAZEL NUT | TUNG |
| HICKORY | WAHOO (Burning Bush) |
| HONEYSUCKLE | WILLOW (Black, Bright) |
| —More Tolerant— | BITTERSWEET |
| ASH (White) | ROSE (Prairie) |
| CEDAR (Red) | |

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, wash off promptly with soap and water. Do not use around the home, recreation areas, or similar areas. Avoid spray drift to susceptible plants—beans, peas, ornamentals, etc. Coarse sprays are less likely to drift. Do not apply directly to flowers, vegetables, fruit trees, ornamentals, cotton or other susceptible plants. This product should be used with caution in clover pastures and a very small amount may injure young grass seedlings. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not contaminate irrigation ditches or water used for domestic purposes. Do not store near fertilizers, seeds, insecticide or fungicides. Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter. Thoroughly clean spraying equipment with suitable chemical cleaner before using for other purposes, or, better yet, do not use the same equipment for other spraying purposes. May cause injury to bentgrass and St. Augustine grass. This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy by burying in a safe place. Do not burn. Do not use, pour, spill or store near heat or open flame.



Contains 1 lb. 2,4-D acid equivalent plus 1 lb. 2,4,5-T acid equivalent per gallon.


ACTIVE INGREDIENTS:
 2,4-dichlorophenoxyacetic acid butyl ester ...
 2,4,5-trichlorophenoxyacetic acid isopentyl ester ...

INERT INGREDIENTS:
 Total
 (equivalent to 2,4-dichlorophenoxyacetic acid 11.9%)
 (equivalent to 2,4,5-trichlorophenoxyacetic acid 11.9%)

CAUTION: Keep Out of Reach of Children.

See other cautions on side panel.
 EPA Est. 159-Ks 1
 E.P.A. Reg. No. 2169-6.

GREEN UP



Brush killer can be dissolved in water. Application can be made until three weeks before cutting, and stump treatment such plants as are listed on forest lands, rights-of-way

DILUTION TABLE
 BRUSH KILLER
 3 to 4 1/2 Pints.
 3 to 4 Quarts.
 1 1/2 to 2 Gallons
 For knapsack and 5 tablespoonfuls of water.

Wet all stems and leaves if necessary for uniform control. If properly applied control 75% or more of brush under some conditions necessary to control regrowth, year is sufficient. If brush be covered to a second year control those missed, or an

STUMP AND TRUNK TREATMENT: Requires 3 gallons of brush killer over the entire area. Make cut stumps and trunks any

NOTICE: Buyer assumes all responsibility for the safe use and/or handling of this product in accordance with the terms of this label.

THIS CONTAINER MAKES

Content

ALL SPECIES OF BRUSH

- Y (Poison)
- NEBERRY
- APPLE (Soft) (Sugar)
- BLACK (Black Jack, Burr, Ash, White)
- POST (White)
- RANGE (Orange)
- RISIMMON
- UM (Sand)
- MAC (Smooth, Staghorn)
- LUMPET VINE
- JING
- AHOO (Burning Bush)
- WILLOW (Black, Bright)
- BRANT
- BITTERSWEET
- DSE (Prairie)

brushed. Avoid contact with skin. In case of contact, wash off with water. Do not use around the house.

Susceptible plants—beans, peas, corn, etc. are less likely to drift. Do not use on fruit trees, ornamentals, or susceptible plants. This product is not for use in clover pastures. It may injure young grass seedlings. Do not use on susceptible plants. Do not contaminate water. Do not use on treated areas. Do not graze meat animals. Do not use for slaughter. Do not use on equipment with suitable for other purposes, or same equipment for other purposes. Do not use on injury to benthos and fish.

Keep out of lakes, streams, rivers, etc. where runoff is likely to occur. Do not use in conditions favoring drift from treated areas. Clean up spilled water by cleaning of spill. Apply this product only to brush.

Dispose of this product by burying in a safe place. Do not pour, spill or store near



green UP




Contains 1 lb. 2,4-D acid equivalent plus 1 lb. 2,4,5-T acid equivalent per gallon.

ACTIVE INGREDIENTS:
 2,4-dichlorophenoxyacetic acid butyl ester
 2,4,5-trichlorophenoxyacetic acid isooctyl ester
INERT INGREDIENTS:
 Total
 equivalent to 2,4-dichlorophenoxyacetic acid (100%)
 equivalent to 2,4,5-trichlorophenoxyacetic acid (100%)

CAUTION: Keep Out of Reach of Children.

See other cautions on side panel.
 E.P.A. Est. 159-K-1
 E.P.A. Reg. No. 2169-6.

GREEN UP



DIRECTIONS

Brush Killer can be dissolved in fuel oil or emulsified with water. Application can be made any time after the full leaf stage until three weeks before frost. Best results are obtained when the brush to be sprayed is under five feet tall. Brush over five feet tall requires repeat application or cutting, and stump treatment. Use along fence rows on such plants as are listed on this label. Do not apply to forest lands, rights-of-way, or pastures.

DILUTION TABLE (For Power Sprayers)

BRUSH KILLER	Water
3 to 4 1/2 Pints	50 Gal.
3 to 4 Quarts	50 Gal.
1 1/2 to 2 Gallons	100 Gal.

For knapsack and hose-attached sprayers use 5 tablespoons of Brush Killer per gallon of water.

Wet all stems and leaves thoroughly as complete coverage is necessary for uniform kill. Amount of material required will depend upon the height and denseness of the brush. If properly applied one thorough application will control 75% or more of the susceptible growth. While under some conditions repeat applications may be necessary to control regrowth, usually a single treatment per year is sufficient. If brush is dense some plants may not be covered so a second application may be necessary to control those missed, or any regrowth.

STUMP AND TRUNK TREATMENT—Dilute at rate of 1/2 pint to 3 gallons of fuel oil. Apply by spraying or brushing over the entire stump or from 18 inches to 6 feet on trunks. Make application heavy until run off is evident on entire area and on all sides. Treat freshly cut stumps and trunks any time of year.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

THIS CONTAINER MAKES UP TO 20 GALLONS OF SPRAY

Contents 1 Quart