DIRECTIONS FOR USE

CRABGRASS CONTROL: Apply 3 pints in sufficient water to cover 1000 sq. ft. of lawn or yard to suppress crabgrass seedlings as germination or sprouting occurs. Use as an early season preventive treatment—not effective for control of growing crabgrass plants
Timing of application is important and should be applied before any germination, which in many areas is during the period April 15 to May 1. This treatment will not burn or injure established lawns. CAUTION: Keep children and pets off treated areas until this material has been thoroughly washed into the soil and the grass has completely dried. Do not apply to newly seeded lawns

CAUTION: Harmful if swallowed. Keep out of reach of children. Avoid contamination of feed and feed stuffs.

Contact with skin can cause toxic symptoms. Avoid spillage on hands and clothing. If spillage occurs, wash with soap and water. Avoid breathing spray mist. Do not use on dairy or meat animals, or in dairy barns and poultry houses. Do not use on feed or forage to be fed to dairy animals or to animals being finished for slaughter.

Toxic to fish and wildlife. Do not contaminate ornamental fish ponds. Note: Birds feeding in treated areas may be harmed.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

45% CHLORDANE

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:	
*Technical Chlordane	45%
Aliphatic Petroleum Hydrocarbons	50%
INERT INGREDIENTS	5%
	100%

*Equivalent to 27% Octachloro-4,7-Methanotetrahydroindane and 18% of related compounds.

CAUTION: Keep out of reach of children.

See other cautions on side panel.

USDA Reg. No. 2169-77

DIRECTIONS FOR USE

FOR LAWN INSECTS	AMOUNT TO USE	DIRECTIONS FOR APPLYING
Grasshoppers Chiggers Wood Ticks Army Worms	2½ oz. (5 tablespoonsful) per gallon of water	Apply for thorough coverage. Repeat as needed. For chiggers and ticks, thoroughly treat lawns, weed patches and other infested areas using 1 gallon diluted spray for each 1,000 sq. ft.
Wireworms White Grubs White Fringed Beetle	4 oz. (8 tablespoonsful) per gallon of water	Apply thoroughly to area, washing well into soil. Use 1 gallon of diluted spray for each 1,000 sq. ft. Repeat as needed.
HOUSEHOLD INSECTS Roaches Waterbugs Silverfish Spiders Ants	6 oz. (12 tablespoonsful) per gallon of water	Spot treat infested areas. Pay particular attention to cracks and crevices where insects may hide or have been observed. Spray along ant trails. Pour one cupful diluted spray into ant nests outside. Repeat as needed to maintain control.

NET WEIGHT 4 OZS