

GENERAL INFORMATION

This product may be used for the control of many broad leaved annual weeds and certain perennial weeds growing in lawns and other turf and in vacant lots and fence rows. In general, this product is most effective when weeds are small and growth is active. It may be applied when the temperatures range from 65° to 90° F. Results obtained above and below these temperatures will be considerably slower and more erratic. Do not apply if rain is expected within 4 hours. Application may be made with either power or hand type sprayers.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, wash off promptly with plenty of water. Avoid spray drift to susceptible plants, as this product may injure cotton, beans, cucumbers, grapes, roses, tomatoes and other vegetable, flower and ornamental plants. Coarse sprays applied at low pressure are less likely to drift. **DON'T OVERDOSE.** Do not store near fertilizers, seeds, insecticides or fungicides. Do not use equipment used in applying this material to apply insecticides or fungicides to susceptible plants or seeds. Do not use on grass seed crops or on turf containing bent grasses. Do not apply to St. Augustine, carpetgrass or clover lawns. Do not use on new grass seedings less than two months old. Clovers present in turf may be damaged. Do not contaminate water used for irrigation or domestic purposes. Do not reuse empty container. Wash thoroughly with water containing detergent. Discard in a safe place.

**THIS CONTAINER WILL MAKE UP TO
128 GALLONS OF SPRAY
NET CONTENTS 1 QUART**

PATTERSON'S®

2,4-D AMINE WEED KILLER

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* 49.4%

INERT INGREDIENTS 50.6%

*2,4-Dichlorophenoxyacetic Acid
Equivalent 41.0%

CAUTION:

Keep out of reach of children.
See other cautions on side panel.

MANUFACTURED BY OR TO THE SPECIFICATIONS OF

PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE

KANSAS CITY, MISSOURI

LABOR SAVING CHEMICALS

6/16/75

54

INFORMATION

and for the control of many
us and certain perennial weeds
other turf and in vacant lots
at this product is most effective
and growth is active. If
the temperature range from
stained above and below these
considerably slower and more er
is expected within 4 hours
with either power or hand

allowed. Avoid contact with
case of contact, wash off
water.
able plants, as this product
cucumbers, grapes, roses
cattle, flower and ornamental
applied at low pressure are
IT OVERDOSE. Do not store
icides or fungicides. Do
n applying this material to
icides to susceptible plants
grass seed crops or on turf
do not apply to St. August
lawn. Do not use on new
two months old. Clovers
aged.

After used for irrigation or
other. Wash thoroughly with
water. Discard in a safe place.

WILL MAKE UP TO
ONS OF SPRAY
NETS 1 QUART

PATTERSON'S

2,4-D AMINE
WEED KILLER

ACTIVE INGREDIENTS

Dimethylamine Salt of 2,4-Dichloro-
phenoxyacetic Acid* 49.4%
INERT INGREDIENTS 50.6%
*2,4-Dichlorophenoxyacetic Acid
Equivalent 41.0%

CAUTION:

Keep out of reach of children.
See other cautions on side panel.

MANUFACTURED BY OR TO THE SPECIFICATIONS OF

PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE KANSAS CITY, MISSOURI

DIRECTIONS

LAWN AND TURF: For control of susceptible or easy-
to-kill weeds in lawn and turf, use 1/2 tablespoon of
this product per gallon of water (1 gallon will cover
approximately 300 square feet) and apply either as
a spot or over-all spray. For hard-to-kill or resistant
weeds, use 1 tablespoon per gallon of water and
apply as above. Repeat treatment as necessary for
control.

VACANT LOTS AND WASTE AREAS: For controlling
broadleaved weed growth in these areas, use 1 table-
spoon of this product per gallon of water and apply
to thoroughly wet all weeds to be killed. Repeat ap-
plication every 2 or 3 weeks as necessary to control
new growth and any regrowth.

WEEDS CONTROLLED

Susceptible or easy-to-kill weeds that are generally
killed with one application:

- Beggar-ticks	- Plantain	- Shepherd's purse
- Cocklebur	- Ragweeds	- Wild buckwheat
- Dandelion	- Sunflower	- Poppygrass
- Horseweed	- Morning-glory	
- Mustard	- Ornamental	

Slightly resistant weeds that require repeated appli-
cation and stronger dosages for control:

- Wild carrot	- Red sorrel	- Curly dock
- Wild garlic	- Sour dock	- Lamb's quarter
- Wild onion	- Big yucca	- Mexican sandbar
- Figweed	- (field)	

NOTICE: Buyer assumes all risk and liability of use,
storage and/or handling of this material not in ac-
cordance with the terms of this label.

LABOR SAVING CHEMICALS

51