

DIRECTIONS FOR USE (Cont.)

into planted fields, and on such places as ditch banks, fence rows and similar non-crop areas. It is best used where there is sparse vegetation under hot dry conditions and immediate contact kill is desired or in dense vegetation where little migration occurs. In the early part of the season when the hoppers are small and immature it should be used as a spray at the rate of 2 1/2 pints concentrate per acre in water sufficient to give good coverage. Late in the season when hoppers are mature and the vegetation is tall and dense the amount of concentrate should be increased to 2 quarts per acre.

PROTECT FROM FREEZING

WARNINGS: Harmful if swallowed. Absorbed through skin. May cause irritation of skin and eyes. Avoid contact with skin, eyes or clothing. In case of skin contact, wash with plenty of soap and water. For eyes, flush with water and get prompt medical attention. Do not breathe vapors or mist from sprays.

Do not use on household pets or humans. Avoid contamination of feeds or foodstuffs. Forage crops treated with this material should not be fed to dairy animals or to animals being finished for slaughter. Do not store near products which will absorb odors. Combustible mixture - Keep container closed and away from heat and open flames.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

NOTICE: Buyer assumes all risk and liability of use, storage, and/or handling of this material not in accordance with the terms of this label.

1 LB. Gamma Isomer Per Gallon

PATTERSON'S

BHC EMULSION CONCENTRATE

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride . . . 11.0%

Other Isomers of Benzene Hexachloride . . . 12.4%

Petroleum Solvent . . . 72.6%

INERT INGREDIENTS . . . 4.0%

TOTAL . . . 100.0% ✓

WARNING: Keep Out of Reach of Children.

See other cautions on side panel.

MANUFACTURED BY OR TO THE SPECIFICATIONS OF

PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE

KANSAS CITY, MISSOURI

USDA Reg. No. 2169-87

APHIDS: To control water (2 teaspoon necessary) - but do

after the fruit starts SPITTLE BUGS: Spr

berries use 1 1/2 pi

ful per gallon of Do not apply mo

CABBAGE MAGG sprouts, cabbage,

mustard greens, c trate mixed with

application should the plants at the

machines vary in plant, the amount

of the machine. pensed with each

in the row and gallons would be

100 gallons, each or the equivalent

BHC Emulsion Co transplanting sta

For use in small mix 1 ounce or 2

and apply 1 pi transplanting.

GRASSHOPPERS: insecticide which

used only on hot

CONTE

Site - Nancy

62

LABOR SAVING CHEMICALS

AC
3/15/71
2109-11

FOR USE (Cont.)

uch places as ditch banks, fence
eas. It is best used where there
ot dry conditions and immediate
n dense vegetation where little
ly part of the season when the
ature it should be used as a
nts concentrate per acre in water
verage. Late in the season when
the vegetation is tall and dense
should be increased to 2 quart

FROM FREEZING

allowed. Absorbed through skin.
n and eyes. Avoid contact with
ase of skin contact, wash with
for eyes, flush with water and
ion. Do not breathe vapors or

cts or humans. Avoid contamina-
Forage crops treated with this
to dairy animals or to animals
r. Do not store near products
ncombustible mixture—Keep container
and open flames
and wildlife. Keep out of lakes,
apply to any area not specified

iner. Destroy by perforating or
safe place.
and liability of use, storage,
not in accordance with

omer Per Gallon

PATTERSON'S

BHC EMULSION CONCENTRATE

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride	11.0%
Other Isomers of Benzene Hexachloride	12.4%
Petroleum Solvent	72.6%
INERT INGREDIENTS	4.0%

TOTAL 100.0%

WARNING: Keep Out of Reach of Children.
See other cautions on side panel.

MANUFACTURED BY, OR TO THE SPECIFICATIONS OF
PATTERSON CHEMICAL COMPANY
1400 UNION AVENUE • KANSAS CITY, MISSOURI

USDA Reg. No. 2169-B7

DIRECTIONS FOR USE

APHIDS: To control in apples use 2 pints per 100 gallons of water (2 teaspoonfuls per gallon of water). ~~Repeat as necessary, but~~ do not use within 60 days of harvest or after the fruit starts to form.

SPITTLE BUGS, STRAWBERRY WEEVIL: For control in strawberries use 1 1/2 pints per 100 gallons of water (1 1/2 teaspoonful per gallon of water). Apply only before blossoms open. Do not apply more than 2 gallons total product per acre.

CABBAGE MAGGOTS: For control in broccoli, Brussels sprouts, cabbage, cauliflower, collards, kale, kohlrabi and mustard greens, apply 1 1/2 gallons of BHC Emulsion Concentrate mixed with sufficient water to cover each acre. This application should be made to the soil around the base of the plants at the time of transplanting. Since transplanting machines vary in the quantity of liquid dispensed to each plant, the amount of water should conform to the capacity of the machine. (If, for example, 1/2 pint of solution is dispensed with each plant, with the plant spacing at 12 inches in the row and 3 feet between rows, approximately 200 gallons would be used per acre. If the machine tank holds 100 gallons, each tankful should be charged with 22 ounces, or the equivalent of 44 tablespoonfuls, per tank.)

BHC Emulsion Concentrate should not be applied after the transplanting stage.

For use in small home gardens on the above named crops, mix 1 ounce or 2 tablespoonfuls, in each 5 gallons of water and apply 1/2 pint of mixture per plant at the time of transplanting.

GRASSHOPPERS: BHC Emulsion Concentrate is a fast killing insecticide which has 3 or 4 days residual effect. It should be used only on hatching areas before the young hoppers move

CONTENTS

GALLON

LABOR SAVING CHEMICALS

ACCT 71

3/15/71

2169-B7

DIRECTIONS FOR USE (Cont.)

into planted fields, and on such places as ditch banks, fence rows and similar non-crop areas. It is best used where there is sparse vegetation under hot dry conditions and immediate contact kill is desired or in dense vegetation where little migration occurs. In the early part of the season when the hoppers are small and immature it should be used as a spray at the rate of 2½ pints concentrate per acre in water sufficient to give good coverage. Late in the season when hoppers are mature and the vegetation is tall and dense the amount of concentrate should be increased to 2 quarts per acre.

PROTECT FROM FREEZING

WARNINGS: Harmful if swallowed. Absorbed through skin. May cause irritation of skin and eyes. Avoid contact with skin, eyes or clothing. In case of skin contact, wash with plenty of soap and water. For eyes, flush with water and get prompt medical attention. Do not breathe vapors or mist from sprays.

Do not use on household pets or humans. Avoid contamination of feeds or foodstuffs. Forage crops treated with this material should not be fed to dairy animals or to animals being finished for slaughter. Do not store near products which will absorb odors. Combustible mixture—Keep container closed and away from heat and open flames.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

NOTICE: Buyer assumes all risk and liability of use, storage, and/or handling of this material not in accordance with the terms of this label.

1 LB. Gamma Isomer Per Gallon

PATTERSON

**BHC EMULSION
CONCENTRATE**

ACTIVE INGREDIENTS:
Gamma Isomer of Benzene
Other Isomers of Benzene
Petroleum Solvent
INERT INGREDIENTS

TOTAL

WARNING: Keep Out of Reach of Children
See other cautions on label

MANUFACTURED BY, OR FOR

PATTERSON CHEMICAL CO.
1400 UNION AVENUE

USDA Reg. No.

LABOR SAVING

3/15/71

2109