

ACCEPTED

DEC 7 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 2164-5 SUBJECT
TO ATTACHED COMMENTS.

DIRECTIONS (CONT'D)

Do not apply to crops later than the number of days appearing in parentheses immediately after the crop name.

CANTALOUPE, WATERMELONS, CUCUMBERS, SQUASH, PUMPKIN, HONEY DEW MELON (5): Melon Worm, Striped Cucumber Beetle, Spotted Cucumber Beetle, Flea Beetle, Pickleworm, Squash Bug, Leafhopper, Melon Aphid, Bacterial Wilt, Angular Leaf Spot and Downy Mildew.

PEPPERS, EGGPLANT, TOMATOES (5): Colorado Potato Beetle, Aphids, Leafhoppers, Flea Beetle, Thrips, Psyllid, Tomato Fruitworm, Tomato Hornworm, Stink Bug, Lace Bug, Lygus Bug, Anthracnose and Cercospora on Peppers, Early and Late Blight on Tomatoes, Anthracnose and Leaf Spot on Tomatoes.

CAUTION: Remove residues remaining at harvest by washing, wiping or similar effective means. Harmful if swallowed or inhaled. Avoid inhalation of dust. Avoid contact with skin. Wash thoroughly after using. Avoid storage near feed and foodstuffs. Keep out of the reach of children.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not reuse container. Destroy when empty.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.



GARDEN DUST

ACTIVE INGREDIENTS:

Zineb (zinc ethylene bisdithiocarbamate)	3.9%
Captan (N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	2.5%
Carbaryl (1-Naphthyl N-methylcarbamate)	2.0%
Rotenone	1.0%
Other cube extractives	1.6%
INERT INGREDIENTS	89.0%

100.0%

INSECTICIDE FUNGICIDE

Ready to Use

CAUTION: Keep out of reach of children.

SEE OTHER CAUTIONS ON BACK PANEL.

Net Contents 1 Pound

EPA Reg. No. 2149-90

DIRECTIONS

Apply GARDEN DUST thoroughly, covering both sides of leaves and the stems as well as possible. Begin applications before diseases are expected to appear and repeat at 7 to 10 day intervals.

Do not apply to crops later than the number of days appearing in parentheses immediately after the crop name. Dust on wet or dry foliage, but preferably when air is still.

GARDEN DUST will control insects and plant diseases listed below:

BEANS (7): Aphids, Japanese Beetle, Leafhopper, Mexican Bean Beetle, Bean Leaf Beetle, Thrips, Downy Mildew.

BEETS (14), CARROTS (7), LETTUCE (14), SPINACH (14): Web Worm, Flea Beetle, Leafhopper, Harlequin Bug, Lygus and Stink Bugs, Leaf Tiers, Alternaria and Cercospora Leaf Spot of Beets and Carrots, Downy Mildew of Lettuce and Spinach.

BLACKBERRIES (14), RASPBERRIES (14): Japanese Beetle, Leaf Roller, Leafhopper, Aphids, Spittlebug, Anthracnose.

STRAWBERRIES (7): Leaf Roller, Meadow Spittlebug, Strawberry Weevil, Japanese Beetle, Botrytis Fruit Rot, Leaf Spot.

GRAPES (7): Leaf Folder, Leafhopper, Leaf Skeletonizer, Berry Moth, Japanese Beetle, Red-Banded Leaf Roller, Downy Mildew, Black Rot.

(Continued Other Panel)