

GENERAL INFORMATION

This product may be used for the control of many broad leaved annual weeds and certain perennial weeds growing in lawns and other turf and in vacant lots and fence rows. In general, this product is most effective when weeds are small and growth is active. It may be applied when the temperatures range from 65° to 90° F. Results obtained above and below these temperatures will be considerably slower and more erratic. Do not apply if rain is expected within 4 hours. Application may be made with either power or hand type sprayers.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, wash off promptly with plenty of water.

Avoid spray drift to susceptible plants, as this product may injure cotton, beans, cucumbers, grapes, roses, tomatoes and other vegetable, flower and ornamental plants. Coarse sprays applied at low pressure are less likely to drift. **DON'T OVERDOSE.** Do not store near fertilizers, seeds, insecticides or fungicides. Do not use equipment used in applying this material to apply insecticides or fungicides to susceptible plants or seeds. Do not use on grass seed crops or on turf containing bent grasses. Do not apply to St. Augustine carpetgrass or clover lawns. Do not use on new grass seedings less than two months old. Clovers present in turf may be damaged.

Do not contaminate water used for irrigation or domestic purposes.

Do not reuse empty container. Wash thoroughly with water containing detergent. Discard in a safe place.

**THIS CONTAINER WILL MAKE UP TO
128 GALLONS OF SPRAY**

NET CONTENTS 1 QUART

PATTERSON'S®

2,4-D AMINE WEED KILLER

ACTIVE INGREDIENTS

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* 49.4%
INERT INGREDIENTS 50.6%
*2,4-Dichlorophenoxyacetic Acid
Equivalent 41.0%

CAUTION:

**Keep out of reach of children.
See other cautions on side panel.**

MANUFACTURED BY OR FOR THE PATTERSON COMPANY

PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE, ST. LOUIS, MISSOURI 63104

DIRECTIONS

LAWN AND TURF: For control of susceptible or easy-to-kill weeds in lawn and turf, use 1/2 tablespoon of this product per gallon of water (1 gallon will cover approximately 300 square feet) and apply either as a spot or over-all spray. For hard-to-kill or resistant weeds, use 1 tablespoon per gallon of water and apply as above. Repeat treatment as necessary for control.

VACANT LOTS AND WASTE AREAS: For controlling broadleaved weed growth in these areas, use 1 tablespoon of this product per gallon of water and apply to thoroughly wet all weeds to be killed. Repeat application every 2 or 3 weeks as necessary to control new growth and any regrowth.

WEEDS CONTROLLED

Susceptible or easy-to-kill weeds that are generally killed with one application:

Beggarticks	Plantain	Shepherd's purse
Cocklebur	Ragweeds	Wild buckwheat
Dandelion	Sunflower	Peppergrass
Horsetail	Morning-glory	
Mustard	Annual	

Slightly resistant weeds that require repeated application and stronger dosages for control:

Wild carrot	Red sorrel	Curly dock
Wild garlic	Sour dock	Lambs' quarter
Wild onion	Bindweed	Mexican sandbar
Pigweed	(field)	

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

E.P.A. Reg. No. 21691-8

LABOR SAVING CHEMICALS