

GARDEN AND TRUCK CROPS

INSECT	CROP	APPLICATION
Tomato Hornworm	Tomatoes*	2½ lbs in 100 gal water per acre
Onion Thrips	Onions*	2½ lbs in 100 gal water per acre, with application repeated if needed
Cabbage Loopers	Cabbage*	1½ lbs in 5 gal water per acre
Cabbage Worms Imported Cabbage Worm		
Harlequin Cabbage Bug	Cabbage	2½ lbs in 100 gal water per acre
Colorado Potato Beetle	Potatoes	2½ lbs in 100 gal water per acre
Beet Webworm	Beets*	2½ lbs in sufficient water to cover 1 acre. Apply early before damage is severe.

*For Garden rate, use 3 gallons of water per 1000 square feet and add 1 tablespoon for each pound shown in the per acre chart

*Do not apply to cabbage, after edible parts begin to form. Do not apply to tomatoes after fruit begins to form. Do not apply to green or spring onions. No application should be made to beets after seedling stage if tops are to be used for food

CRABGRASS CONTROL

Apply 3½ to 4 lbs to each 1000 square feet of lawn to suppress crabgrass seedlings as they germinate. Use as an early season preventative treatment not effective for control of growing crabgrass plants. Timing of application is important and should be applied before any germination occurs, which in many areas is during the period April 15th to May 1. This treatment will not burn or injure established lawn. Do not use on newly seeded lawns. Keep children and pets off treated areas until this material has been completely watered into the soil and the grass has completely dried. Do not contaminate irrigation ditches or water used for domestic purposes. This material can also be applied as a liquid spray with water 4 lbs of 40% Chlordane in 10 to 15 gallons of water for 1000 sq ft area.

CAUTION: Harmful if swallowed. Contact with skin causes toxic symptoms. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Avoid breathing dust or spray mist. Keep out of reach of children. Do not apply to dairy animals, calves or in dairy barns. Do not use on feed or forage crops to be fed to dairy animals or those being finished for slaughter. Do not contaminate irrigation ditches or water used for domestic purposes.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift of spray from areas treated. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Container should be disposed of by punching holes in them and burying them with wastes or by burning. Keep out of smoke.

NOTICE: Buyer assumes all risk and liability of use, storage and or handling of this material not in accordance with the terms of this label.

PATTERSON'S®

40% CHLORDANE

WETTABLE POWDER

ACTIVE INGREDIENTS:
*Technical Chlordane 40.0%

INERT INGREDIENTS 60.0%

TOTAL 100.0%

*Equivalent to 24% Octachloro-4,7-Methanotetrahydroindane and 16% related compounds.

CAUTION:
Keep Out of Reach of Children.
See back panels for additional warnings.

NET CONTENTS 1 POUND

PATTERSON CHEMICAL COMPANY

USDA Reg No 2169 99

DIRECTIONS FOR USE

SOIL INSECTS ATTACKING CORN, BEETS, BEANS, CABBAGE AND LAWNS
Unless otherwise specified, rates are for dilution with sufficient water to cover one acre.

INSECT	RATE OF APPLICATION	TIME OF APPLICATION
Ants	1½ lbs per 3 gallons of water 2% solution	Thoroughly wet soil or turf at first sign of infestation
Wireworms	6½ lbs	Work into soil to 4-6" depth at planting time
White Grubs	12½ 25 lbs	Apply to soil and turf in Spring
White-Fringed Beetle Larva	6½ lbs	Work into soil to 4-6" depth at planting time
Asiatic Garden Beetle	25 lbs	Work into soil to 4-6" depth at planting time
Cabbage Maggots	2½ 5 lbs	Work into soil to 4-6" depth at planting time
Chinch Bugs	4 tbs in 5 gals of water per 1000 sq ft	Apply to soil surface at first signs of infestation. Mow lawn, remove thatch, water thoroughly, apply Chlordane and water in. Repeat within 10 days.
Onion Maggots	2½ lbs	Apply when seeding
Catworms	8 tbs in 5 gals of water per 1000 sq ft	Apply to soil surface around plants
Corn Rootworm	2½ lbs	Apply when seeding as row treatment
Seed Corn Maggot	2½ 5 lbs	Apply when seeding
Mole Crickets	8 tbs in 10 gals of water per 1000 sq ft	Apply to soil surface around plants. Water area before application
Sod Webworm Lawn Moth	1 lb per 20 gals of water	Apply at 6 to 7 gallons per 1000 sq ft
Chiggers	8 tbs in 5 gals of water per 1000 sq ft	Keep children and pets off treated lawn areas until insecticide has been washed into the soil and the grass is thoroughly dry.

Do not use on newly seeded lawns.

RESIDUAL CONTROL

Prepare a spray by diluting 1 pound with 2½ gallons of water. Apply this spray as a spot treatment to infested areas such as cracks, crevices, the undersides of carpets and infested areas of top surface for the control of carpet beetles, cockroaches, scorpions, silverfish, spiders, black widow included.

LABOR SAVING CHEMICALS

10-11-1968
1169 99