

READ DIRECTIONS AND CAUTION BEFORE USING TO BE USED ON THE FOLLOWING CROPS:
 Alfalfa and Clover for Forage, Celery, Cucurbits, Spinach, also Greenhouse Nursery Plants, Camellias, Azaleas and Rhododendrons.

Directions

Alfalfa Weevil: Use ¾ pint per acre in sufficient water for even distribution. Apply when larvae feeding damage begins to show in the field.

Lygus Bugs, Thrips: 1 pint per acre in sufficient water for even distribution.

Pea Aphid: 1½ to 2 pints per acre in sufficient water for even distribution.

Note: On above do not use later than 2 weeks before cutting or pasturing and do not use more than ½ pound actual lindane per acre.

Celery: For Aphid, Thrips, Tarnished Plant Bugs, and certain types of Lepidopterous Larvae use 1 pint per 100 gals. water.

Note: Do not apply after plants are half grown.

Cucurbits: For Cucumber Beetles, Thrips use ¼ to ½ pint per 100 gals. water.

Note: Do not use after blossoms form.

Squash Bugs: On Squash use 1 to 2 pints per 100 gals. water.

Note: Do not use after blossoms form.

Aphid: Use 1 pint per 100 gals. water.

Note: Do not use after blossoms form.

Spinach: For Aphid, Leaf Miners use 1 pint per 100 gals. water.

Note: Apply only in seedling stages.

Greenhouse Nursery Plants: For Aphid, Thrips, Loopers, Leafrollers, Plant Bugs, White Fly, Psyllid, Leaf Miners, Flea Beetles, Diabrotica Beetles use 1 pint per 100 gals. water. Repeat application as necessary.

EACH GALLON MAKES UP TO 3200 GALLONS OF SPRAY

NET CONTENTS _____ GALLON

PATTERSON'S

20% LINDANE

EMULSIFIABLE CONC.

ACTIVE INGREDIENTS:

*Gamma Isomer of Benzene Hexachloride from Lindane
 Petroleum Solvents
 INERT INGREDIENTS

BY WEIGHT

20%
 72%
 8%
 100%

*Lindane per gal.—1.65 pounds.
 Lindane No. 20 is a liquid formulation containing Lindane. It forms an emulsion with water and is suitable for use in conventional spraying equipment including Airplane, Fog Machines, and Aerosol equipment.

CAUTION: Keep out of reach of children.

MANUFACTURED BY PATTERSON CHEMICAL COMPANY

PATTERSON CHEMICAL CO.

1400 UNION AVENUE • KANSAS CITY, MO.

Y
 RI

Directions

5 Camellias, Azaleas, Rhododendrons: For Fuller Ros Brachyrhinus, Cucumber Beetle Larvae on Roots use per 100 gals. water. Spray to saturate soil around in beds.

8 For potted plants: Spray soil surfaces lightly, cool to a light rain.

9 Soil Insects: Wire worms, Cutworms, White Grub, Brötica Larvae and certain other soil insects. Use pints per acre in sufficient water for even dist. Apply as a spray directly on the soil, then disc as practicable to give thorough distribution in the top 6 inches of soil. Do not use in soil to be planted crops.

Termites: (Subterranean) Use 2 gallons 20% Lindane to 50 gals. of water or oil. Apply by spraying into shallow trenches around building foundation below surface. Apply 2 gallon per 5 lineal feet of shallow and 4 gallon per 5 lineal feet in trench 30 inches deep. Soil surface, wooden studs, etc., under building also be treated in case of severe infestations. surface treatment, apply 1 gal. diluted spray per 14 Make annual inspections for possible reinfestation

CAUTION: Various restrictions have been placed use of pest control chemicals on food crops. Avoid of this product on edible portions of food crop there is a possibility of residue remaining at harvest contamination of feed and foodstuffs and avoid in contact with skin and eyes may cause irritation. absorbed through skin. Wash with plenty of water. For eyes, get medical attention—flush with for fifteen minutes. Treated forage crops, other Alfalfa and Clover, are not to be fed to dairy animals being finished for slaughter.

NOTICE TO BUYERS: Seller makes no warranty kind, express or implied, concerning the use of the product. Buyer assumes all risk of use or handling when accordance with directions or not.

USDA Reg. No. 2169-110

LABOR SAVING CHEMICALS

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT AND REGULATIONS THEREUNDER THIS PRODUCT IS SUBJECT TO ATTACHED COMMENTS.

Check for copy

PATTERSON'S®

20% LINDANE

EMULSIFIABLE CONC.

ACTIVE INGREDIENTS:	BY WEIGHT
*Gamma Isomer of Benzene Hexachloride from Lindane	20%
Petroleum Solvents	72%
INERT INGREDIENTS	8%
	100%

*Lindane per gal.—1.65 pounds.
Lindane No. 20 is a liquid formulation containing Lindane. It forms an emulsion with water and is suitable for use in conventional spraying equipment including Airplane, Fog Machines, and Aerosol equipment.

CAUTION: Keep out of reach of children.

See other cautions on the side of the can.

PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE • KANSAS CITY, MO. 64108

Directions

⁵ ⁶ ⁷ **Camellias, Azaleas, Rhododendrons:** For Fuller Rose Weevil, Brachyrhinus, Cucumber Beetle Larvae on Roots use 1 pint per 100 gals. water. Spray to saturate soil around plants in beds.

⁸ **For potted plants:** Spray soil surfaces lightly, comparable to a light rain.

⁹ **Soil Insects:** Wire worms, Cutworms, White Grubs, Diabrotica Larvae and certain other soil insects. Use 2 to 2½ pints per acre in sufficient water for even distribution. Apply as a spray directly on the soil, then disc as soon as practicable to give thorough distribution in the upper 4 to 6 inches of soil. Do not use in soil to be planted to root crops.

Termites: (Subterranean) Use 2 gallons 20% Lindane Spray to 50 gals. of water or oil. Apply by spraying into narrow, shallow trenches around building foundation below floor surface. Apply 2 gallon per 5 lineal feet of shallow trench and 4 gallon per 5 lineal feet in trench 30 inches or more deep. Soil surface, wooden studs, etc., under buildings may also be treated in case of severe infestations. For soil surface treatment, apply 1 gal. diluted spray per 100 sq. ft. Make annual inspections for possible reinfestation.

CAUTION: Various restrictions have been placed on the use of pest control chemicals on food crops. Avoid the use of this product on edible portions of food crops where there is a possibility of residue remaining at harvest. Avoid contamination of feed and foodstuffs and avoid inhalation. Contact with skin and eyes may cause irritation. May be absorbed through skin. Wash with plenty of soap and water. For eyes, get medical attention—flush with water for fifteen minutes. Treated forage crops, other than Alfalfa and Clover, are not to be fed to dairy animals or to animals being finished for slaughter.

NOTICE TO BUYERS: Seller makes no warranty of any kind, express or implied, concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

USDA Reg. No. 2169-110

LABOR SAVING CHEMICALS

1968
VICIDE
AG
PATERSON