

FRUITS: APPLES, PEARS, CRABAPPLES, QUINCES: For European red mite, clover mite, two-spotted (red spider), Schoene, McDaniel, yellow (carpini), Willamette, Pacific, and apple rust mites. Apply in second or later cover sprays or whenever mites appear but not later than within 7 days of harvest.

PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS: For European red, clover (brown almond), two-spotted (red spider), Schoene, McDaniel, Pacific, Willamette, peach silver, plum nursery and yellow mites.

Apply at petal fall to first cover spray or whenever mites appear. If repeat application is necessary, wait 30 days to avoid excessive residues. Do not apply to cherries, plums, prunes or figs within 7 days of harvest, or on other fruits listed above within 14 days of harvest.

CITRUS FRUITS: For citrus red (purple), citrus flat, six-spotted and Yuma mites, on non-bearing trees or as post-harvest application on bearing trees, follow local timing recommendations obtainable from State Extension Service Specialist.

CAUTION: Harmful if swallowed. Avoid inhalation of spray mist; skin contact, contamination of feed or food-stuffs. Wash thoroughly after handling.

This product is toxic to fish. Keep out of lakes, streams and ponds.


NOTICE: Buyer assumes all risk and liability of use, storage and or handling of this material not in accordance with the terms of this label.

ACTIVE INGREDIENTS:

1,1-Di-p-chlorophenyl	18.5%
2,4,2-trichloroethanol	70.0%
Xylene	11.5%
INERT INGREDIENTS	100.0%

Do not use, pour, spill or store near heat or open flame.

*Kelthane is the registered trademark of
Rohm & Haas Co.



Kills MITES on Roses, Evergreens, Flowers, Vegetables, Fruits, Berries, Nuts, Shade Trees, Ornamentals and Nursery Stock. Kills CLOVER MITES outside homes.

CAUTION:

Keep out of reach of children.
See other cautions on side panel.

EPA Reg No 2169142 8 73

DIRECTIONS

Use 2 teaspoonfuls in 1 gallon of water, applying thoroughly to both upper and under sides of foliage when mites first appear and as necessary thereafter.

FLOWERS, ORNAMENTALS, NURSERY STOCK, TREES: To control two-spotted mite (red spider), European red mite, Spruce mite and other mites common on both greenhouse and outdoor plants such as evergreens, shade trees, flowers, nursery stock and ornamentals, use on Canada juniper.

VEGETABLES: To control two-spotted mite (red spider), Pacific Tropical and tomato russet mites on beans, cucumbers, cantaloupes, melons, watermelons, pumpkins, winter and summer squash, tomatoes, eggplants and peppers. Do not use on beans within 7 days of harvest or on other vegetables within 2 days of harvest. Do not feed vines to meat or dairy animals.

STRAWBERRIES: To control two-spotted mite (red spider), strawberry mite and six-spotted mite, use as above. To control cyclamen mite use 4 to 6 teaspoonfuls per gallon water and drench plants thoroughly at 10 to 20 day intervals in spring when new growth starts. Repeat as necessary. Do not use within 2 days of harvest.

CLOVER MITES OUTSIDE HOMES: To control clover mites, spray walls, foundations, window-sills; also plants and lawn at base of building.

BERRIES, GRAPES, NUTS as listed: To control brown and two-spotted (red spider), European red, Pacific, McDaniel, and yellow mites on grapes, raspberries, dewberries, loganberries, boysenberries, walnuts, filberts, pecans, chestnuts and hickory nuts, apply just before bloom or whenever mites appear. On nuts do not apply within 14 days of harvest. Do not use on grapes within 7 days of harvest, or on berries within 2 days of harvest.

NET CONTENTS ONE PINT