

SEED TREATMENT: BEANS, CABBAGE, CORN, MELONS, PEAS, SQUASH (Seed Rot, Damping-Off)— $\frac{1}{2}$ to 1 teaspoonful per pound of dry seeds. BEETS, CHARD, SPINACH, GRASS— $1\frac{1}{2}$ to 2 teaspoonfuls per pound of dry seed. Mix thoroughly in paper bag or glass jar. Do not use on Lettuce Seed.

FRUIT: APPLE (Scab, Fruit Spot, Sooty Blotch, Fly-Speck, Frog-Eye, Fruit Rots)—8 teaspoonfuls per gallon of water. Apply at about weekly intervals in the pre-bloom, bloom, petal-fall, first and second cover sprays; 10 to 14 day intervals in the third and later cover sprays. PEACHES (Brown Rot, Scab)—8 teaspoonfuls per gallon of water in full pink, bloom, petal-fall, shuck, cover and pre-harvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control Blossom-Blight. Repeat application at 7 to 14 day intervals as needed to maintain cover. CHERRY (Brown Rot, Leaf Spot)—8 teaspoonfuls per gallon of water in full pink, bloom, petal-fall, shuck, cover and pre-harvest sprays. Application at 3 to 4 day intervals may be necessary during bloom to control Blossom-Blight. Repeat application at 7 to 14 day intervals as needed to maintain cover.

STRAWBERRY: (Botrytis Rot)—8 teaspoonfuls per gallon of water. Make first application as soon as new growth starts and repeat at 6 to 10 day intervals to harvest. Under severe infection conditions apply immediately after each picking through harvest.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label

PATTERSON'S®

CAPTAN GARDEN SPRAY

FOR PLANT DISEASE CONTROL

ACTIVE INGREDIENT:

Captan (N-trichloromethyl-
thio-4-cyclohexene-1, 2-
dicarboximide) 50%

INERT INGREDIENTS 50%

CAUTION:

Keep Out of Reach of Children.

Avoid contamination of feed and foodstuffs.
Avoid inhalation of dust or spray mist. Avoid
prolonged or repeated contact with skin.

This product is toxic to fish.
Keep out of lakes, streams and ponds.

NET WEIGHT 1 LB.

PATTERSON CHEMICAL COMPANY
1400 UNION AVENUE • KANSAS CITY, MISSOURI

DIRECTIONS FOR USE

ORNAMENTALS: AZALEA, CARNATION, CHRYSANTHEMUM (Damping-Off)—Dip cuttings in a suspension of 5 tablespoonfuls per gallon of water before bedding. CARNATION (Rust, Alternaria Leaf Spot); CHRYSANTHEMUMS (Botrytis Flower Blight)—8 teaspoonfuls per gallon of water. Apply at first sign of disease. Repeat at 6 to 10 day intervals. Use the shorter interval under conditions favorable to infection. GLADIOLUS, TUBEROUS BEGONIAS (Rot, Damping-Off)—8 tablespoonfuls per gallon of water. Dip bulbs for 30 minutes. Stir, drain and plant. ROSE (Black Spot)—8 teaspoonfuls per gallon of water. Apply at first sign of disease. Repeat at 6 to 10 day intervals. Use the shorter interval under conditions favorable to infection.

VEGETABLES: CANTALOUPE, CUCUMBER, PUMPKIN, SUMMER AND WINTER SQUASH, WATERMELON: (Angular Leaf Spot, Anthracnose, Downy Mildew)—8 teaspoonfuls per gallon of water. Apply at 6 to 10 day intervals. Use the shorter interval under conditions favorable to infection. Begin applications at first sign of infection. POTATO (Early and Late Blight), TOMATO (Early and Late Blight, Grey Leaf Spot, Septoria Leaf Spot)—8 teaspoonfuls per gallon of water. Apply at 6 to 10 day intervals. Use the shorter interval under severe infection conditions. Begin applications at first sign of infection. CELERY (Damping-Off)—8 teaspoonfuls per gallon of water per 100 square feet of seedbed. Make first application when all seeds have germinated and seedlings have taken root. Repeat at 6 to 10 day intervals. Use the shorter interval under conditions favorable to damping-off.

USDA Reg. No. 2169-159

LABOR SAVING CHEMICALS