




FOR FLOWERS AND ORNAMENTALS
VEGETABLES AND TURF



ACTIVE INGREDIENT:
Zineb (zinc ethylene bisdithiocarbamate) . 75%
(Metallic Zinc Equivalent 17.7%)
INERT INGREDIENTS. 25%
100%

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT


(**SWALLOWED:** Call a physician or Poison Control Center immediately. Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.

EPA Reg. No. 2169-160

EPA Est. No. 2169-MO-1

NET CONTENTS _____ (6 Ounces; 12 Ounces)

PATTERSON GREEN-UP COMPANY
Div. Curry Cartwright, Inc.
1331 Union Avenue Box 4029
Kansas City, MO. - 64101



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS &
DOMESTIC ANIMALS**

CAUTION

Harmful if inhaled or swallowed. May cause irritation of eyes, nose, throat and skin. Do not breathe dust or spray mist. In case of skin contact wash with water, for eyes wash with water and get medical attention.

Protective Clothing -- long pants, long sleeved shirt, gloves, hat and boots -- must be worn during mixing and loading of the product.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

CHEMICAL OR PHYSICAL HAZARDS

Keep away from fire and sparks. Do not allow to become wet or overheated during storage. This may bring on chemical changes which will impair the fungicidal effectiveness and may also generate flammable vapors. Keep container closed when not in use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked, place out of reach of children. Do not reuse empty container. Wrap container and put in trash.

Note: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

General Instructions: Use ZINEB FUNGICIDE with water at the rate of 2 to 4 level teaspoonfuls per gallon. Thorough and uniform coverage is absolutely essential for disease control. Start applications before diseases are apparent (for fruit, make first applications just before bloom and just after bloom) and protect plants through the growing season by applying at 5 to 10 day intervals. Unless otherwise indicated, do not apply to any food crop within 7 days of harvest.

DISEASES CONTROLLED

FLOWERS AND ORNAMENTALS

Azaleas and Camellias - For Petal Blight. Spray 2 to 3 times a week while flowers are opening. Direct the spray into the flowers, Spray the ground under the bushes thoroughly.

Carnations and Snapdragons - For Leaf Spots and Rusts.

Chrysanthemums - For Septoria Leaf Spot and Rust.

Cyclamen - For Botrytis Blight and Leaf Spot.

Gladioli - Septoria Leaf Spot, Stemphyllium, Botrytis Blight and Curvularia.

Tulips - Botrytis Blight, (Tulip Fire).

Roses - For control of Black Spot.

BEST AVAILABLE COPY

DISEASES CONTROLLED (Cont.)

VEGETABLES

Tomatoes - For Late and Early Blight, Septoria Leaf Spot, Gray Leaf Spot and Anthracnose. Do not apply within 5 days of harvest.

Melons, Pumpkins, Cucumbers, Squash - For control of Downy Mildew, Alternaria, Gummy Stem Blight, Angular Leaf Spot and Anthracnose. Do not apply within 5 days of harvest.

Beans (Snap) - For control of Rust and Anthracnose.

Beans (Limas) - For control of Anthracnose and Downy Mildew.

Spinach - For Downy Mildew, and White Rust.

Cabbage, Cauliflower, Broccoli - For control of Downy Mildew.

Sweet Corn - For control of Helminthosporium Leaf Blight.

Eggplant - For Alternaria, Anthracnose and Phomopsis.

Peppers - Anthracnose, Cercospora Leaf Spot

Carrot and Beets - Leaf Spots.

FRUIT - Apply at 7 to 14 days intervals beginning before bloom. Thoroughly spray.

Apples - For control of Apple Scab, Apple Rust, Black Rot, Brooks Spot, Sooty Blotch, Fly Speck.

Cherries - For Leaf Spot. Apply at petal fall; 10 to 20 days later and again immediately after harvest.

Grapes - For control of Black Rot and Downy Mildew. Apply just before bloom; just after bloom; 10 days after bloom; and 2 to 3 weeks later.

Pears - Scab and Sooty Blotch.

Strawberries - For control of Leaf Spot. Begin when new growth starts in the spring and repeat as necessary.

BEST AVAILABLE COPY