




FOR ORNAMENTALS & VEGETABLES

ACTIVE INGREDIENT

Maneb (Manganese ethylene
bisdithiocarbamate)

Manganese equivalent as metallic 16.5%

NET INGREDIENTS

Total

80.0%

20.0%

100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

If Swallowed — Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat. Never give anything by mouth to an unconscious person.

If In Eyes — Wash with water for at least 15 minutes. Get medical attention.

If On Skin — Wash thoroughly with soap and water.

EPA Reg. No. 216-161 EPA Est. No. 2159-15-1

Net Contents 8 Ounces

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

VEGETABLES

HOW TO MIX — For the Homeowner — Mix 1 level tablespoonful per gallon of water. Add a Spreader Sticker to the mixture to give thorough coverage of plants. Use a hose-end sprayer or small pressure sprayer suitable for spraying wettable powders. Spray plants to point of run-off. **WHEN TO USE** —

Unless otherwise noted, begin applications when diseases first appear and repeat at 7 to 10 day intervals throughout the growing season in conditions favoring disease development. Do not spray the listed days between the last spray and harvest.

BEANS — Including Limas. For control of Downy Mildew, Rust and Anthracnose. Apply at 4 to 7 day intervals when diseases first appear. Do not apply to string beans within 4 days of harvest.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI — For Alternaria Leaf Spot and Downy Mildew. Use at 3 day intervals in plant beds. Do not apply within 7 days of harvest.

CANTALOUPE, CUCUMBERS, MELONS, SQUASH, WATERMELONS — For Anthracnose, Downy Mildew and Gummy Stem Blight. Begin applications when plants begin to run and under severe disease conditions repeat applications should be shortened to 3 to 5 day intervals. Do not apply within 5 days of harvest.

COLLARDS, KALE, MUSTARD GREENS, TURNIPS — For Alternaria Leaf Spot and Downy Mildew. Remove residues by washing. Do not apply within 10 days of harvest.

EGGPLANT, PEPPERS — For Anthracnose and Fruit Rot (Phomopsis vexans). Apply as fruit begin to form and repeat at weekly intervals through harvest. Remove residues by washing or other effective means.

LETTUCE (heads and leaf) — For Downy Mildew. Apply at first sign of disease and continue as indicated above. Remove residues by stripping from heads and washing. Do not apply within 10 days of harvest.

TOMATOES — For Early and Late Blight, Septoria Leaf Spot, Late Leaf Spot, Stemphylium, Glados, Late Gray Mold and Anthracnose on tomatoes in the field. Do not apply on skin of tomatoes. Treat when diseases first appear in the area and continue at 7 to 10 day intervals in the field. Under severe blight conditions use at 3 day intervals for severe Anthracnose outbreaks. To avoid damage, do not use on tender young plants in glass. Do not apply within 5 days of harvest.

ORNAMENTALS

Use 1 level tablespoonful per gallon of water and spray plants to give full coverage. This is equivalent to 1.6 fluid ounces per 100 gallons of water.

Begin spraying when plants are well-leaved out and disease first appears and repeat at 7 to 10 day intervals throughout the growing season when conditions favor disease development.

CARNATION — For Alternaria Leaf Spot. Use 1 level tablespoonful per gallon of water to give full coverage.

DAHLIA — Botrytis Flower Blight — 1 level tsp. per gallon water.

GLADIOLUS — Leaf and Flower Spot, Botrytis & Curvularia. Respectively.

LILY — Botrytis Blight.

PANSY — Anthracnose. Use same rate as for carnations.

PEONY — For Botrytis Blight and Phytophthora Blight. Apply in early spring and early fall by drenching soil around the plants as well as full coverage on foliage. Destroy infected parts promptly.

ROSES — For Black Spot. Begin applications when leaves first unfold.

SNAPDRAGON — For Rust.

ZINNIA — For Leaf Blight, Alternaria Sp. Use 1 level tsp. per gallon of water. Use a Spreader Sticker to insure proper coverage of Zinnia foliage.

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

CAUTION

Harmful if inhaled or swallowed. May cause irritation of eyes, nose, throat and skin. Do not breathe dust or spray mist. In case of skin contact wash with water. For eyes wash with water and get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

CHEMICAL OR PHYSICAL HAZARDS

Keep away from fire and sparks. Do not allow to become wet or overheated during storage. This may bring on chemical changes which will impair the fungicidal effectiveness and may also generate flammable vapors. Keep container closed when not in use.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry place.

Do not reuse empty bags or canisters. For canister wrap and put in trash collection. For bags dispose of according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

NOTICE — Buyer assumes all risk and liability of use, storage and/or handling of this material in accordance with the terms of this label.

P.O. Box 4029

patterson GREEN UP company

Kansas City, Missouri 64101

Division Curry-Cartwright Inc.

301 858-26

15

1

16

Fatterson Greenup Company
1400 Union Avenue
Kansas City, MO 64101

Attention: Miles R. McIane

Gentlemen:

Subject: Manel Fungicide
EPA Registration No. 2169-161
Your Application for Amended Registration
Dated July 31, 1981

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable, provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. Add a statement to the label requiring protective clothing (long pants, long sleeve shirt, gloves, hat and boots) be worn during mixing and loading.
 - b. Highlight preharvest intervals on labels of noncommercial (homeowner) products.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

01-11
-2-

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure