

**CONTROLS JAPANESE BEETLES, MEXICAN BEAN BEETLES, SQUASH BUGS AND CERTAIN OTHER INSECT PESTS OF FRUIT, VEGETABLES AND ORNAMENTALS.**

Patterson's 50% Sevin\* Wettable Powder is a wettable spray powder. Measure in level tablespoonfuls. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline materials. It is compatible with most fungicides and miticides. Do not mix with oil base sprays.

**ORNAMENTALS:** Use 2 tablespoonfuls per gallon for Japanese beetles, flea beetles, bagworms, leafhoppers, elm leaf beetles, certain aphids, blister beetles, leaf rollers, plant bugs, scale insects, box elder bugs and tent caterpillars. Apply when pests appear. Repeat as needed.

**APPLES AND PEARS:** Use 2 tablespoonfuls per gallon for codling moth, pear psylla, apple maggot, plum curculio, fruit tree leaf rollers, Japanese beetles, apple aphids and San Jose scale. Do not apply before second cover spray to avoid fruit thinning. Apply at 10 day intervals in July for apple maggot. May be used up to 1 day before picking.

**PEACHES AND APRICOTS:** Use 1 tablespoonful per gallon for Oriental fruit moth, plum curculio, peach twig borer, cat-facing bugs, San Jose scale, Japanese beetles. May be used 1 day before harvest on peaches; 3 days on apricots.

**GRAPES:** Use 1 tablespoonful per gallon for leafhoppers, grape berry moth, Japanese beetles. May be applied up to harvest.

**STRAWBERRIES:** Use 2 tablespoonfuls per gallon up to 1 day before picking for leaf roller and meadow spittlebug control. Make 3 to 5 post bloom summer sprays as needed.

**BEANS:** Use 2 tablespoonfuls per gallon. Spray until all foliage is wet. Repeat in 7 to 10 days. For Mexican bean beetles, leafhoppers, flea beetles, stink bugs, lima bean pod borer. May be used up to harvest. Do not feed vines to dairy or meat animals.

**CUCUMBER, MELONS, AND SQUASH:** Use 2 tablespoonfuls per gallon. Apply as full wetting spray for squash bugs, melon worms, cucumber beetles and pickle worms.

**POTATO:** Use 2 tablespoonfuls per gallon for Colorado potato beetle, leafhopper and flea beetles.

**TOMATO, PEPPERS, EGGPLANTS:** Use 2 tablespoonfuls per gallon for tomato fruit worm, flea beetles, Fall army worms, stink bugs and leafhoppers. Apply up to day of picking.

**CAUTION: HARMFUL IF INHALED OR SWALLOWED.** Do not breathe dust. Do not take internally. Avoid breathing of dust. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Do not apply to open blossoms. This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. "This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service." Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**NOTICE:** Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.



**This Package Makes Up to 20 Gallons of Spray**

**ACTIVE INGREDIENT:**

Carbaryl (1-Naphthyl N-Methylcarbamate) 50%

**INERT INGREDIENTS** ..... 50%

**100%**

\*Trade Mark Union Carbide Corporation for 1-Naphthyl N-Methylcarbamate. U.S. Patent 2,903,478.

**CAUTION:**

**Keep Out of Reach of Children.  
See other warnings on back panel.**

E.P.A. Reg. No. 2169-165

**NET WEIGHT 6 OZS.**

E.P.A. Est 2169-Ks.-1

**Patterson**