

BENT GRASSES may be injured with this spray. Do not apply to newly seeded lawns or treat any new desirable grass seedlings until mowed at least twice. Reseeding after treatment should be delayed about one month.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin and clothing. Wash after using.

Avoid spray drift to susceptible plants, as this product will injure grapes, tomatoes, roses, beans, boxelder, zinnias, etc. Do not use the same sprayer for other purposes until cleaned with a suitable cleaner. Do not apply to St. Augustine, Dichondra or carpetgrass lawns. Do not use in greenhouses.

Vapors from this product may harm nearby susceptible plants. When spraying, coarse sprays are less likely to drift. Keep away from heat and open flame. Do not store near fertilizers, seeds, insecticides or fungicides. This product is toxic to fish. Keep out of lakes, streams or ponds.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.



CHICKWEED KILLER

CONTAINING SILVEX

ACTIVE INGREDIENTS:

*2-Ethylhexyl ester of Silvex 2-(2,4,5-trichlorophenoxy) propionic Acid	13.3%
INERT INGREDIENTS	86.7%
	<hr/> 100.0%

*Equivalent to 9.4% of 2-(2,4,5-trichlorophenoxy) propionic Acid (Silvex).

CAUTION: Keep out of reach of children.
See other cautions on side panel.

E.P.A. Reg. No. 2169-166

E.F.A. Est. 2169-Ks.-1

DIRECTIONS FOR USE

For Chickweed, mix 2 tablespoonfuls per 1 gallon of water and apply uniformly over approximately 130 sq. fet. of lawn. Treat early in the fall or early in the spring, or at any time when Chickweed is actively growing. It may be applied at the same rate during the summer months to control White Clover, Black Medic, Veronica, Knotweed, Ground Ivy, as well as Dandelions, Plantains and Buckhorn. Do not use in flower beds or around ornamental plantings. Knotweed should be treated soon after the weed emerges.

SPOT SPRAYING: For small patches, use 1 tablespoonful in 1 gallon water and apply to thoroughly cover the foliage of undesirable weed. Use care not to overdose, but wet thoroughly.

CLOVER IN LAWNS: To stop clover from blooming, but not completely kill it, mix 1 tablespoonful per gallon water and apply to about 220 sq. ft. in March or April.

NET CONTENTS 1 QUART