

Garden Weed Preventer

(CONTAINS DACTHAL*)

**PREVENTS NEW WEED GROWTH IN FLOWER
AND VEGETABLE GARDENS**

One Pound Treats 100 Sq. Ft.



ACCEPTED
2169-180

BEST AVAILABLE COPY

ACTIVE INGREDIENT
Dimethyl Tetrachloroterephthalate**
INERT INGREDIENTS

	25%
	97.5%
TOTAL	100%

*Trade Mark Diamond Shamrock Co.
**U.S. Patent 2,923,634. For use as pre-emergent herbicide.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See Back Panel For Additional Caution Statements

EPA Reg No 2169-180

EPA EST No 2169-KS-1

NET WEIGHT 15 POUNDS



DIRECTIONS FOR USE

LAWNS & ORNAMENTAL TURF — Use 25 lbs. to treat 2500 sq. ft. Apply in the spring (March through April) prior to germination of Crabgrass, Foxtail, and other annual weeds. Does not provide control after weeds germinate. Wait 60 days after application before sowing grass seed to treated areas. Use a good lawn fertilizer spreader at following rates — SCOTT & LAWN BEAUTY - 9; CYCLONE & SIMILAR TYPES - 6.

Read ALL the Label FIRST! To be effective, existing weeds must be removed in early spring or treatment made on freshly cultivated ground. Apply 1 pound per 100 square feet. Sift over ground as uniformly as possible. Water lightly after application. Do not treat dry, hard ground. Do not cultivate ground after application. A second late Summer or Fall application gives year around weed free plantings in the listed ornamentals, shrubs and flowers. See cautions for plants that should NOT be treated.

Controls crabgrass, foxtail, purslane, lambsquarters, red root pigweed, common chick weed, spotted spurge, as well as many other annual grasses and broadleaf weeds. Ragweed and smartweed are not controlled.

May be used around and under shade trees, evergreens, flowering shrubs and hedge rows. Also in ground covers such as ivy and pachysandra. Do not use on dichondra. Does not prevent growth into treated areas by turf grasses.

May be used very effectively in rose beds and in iris and gladiolus plantings.

Treat annual flowers after transplanting from flats or after thinning and initial weeding. Do not treat immediately after seeding, petunias, snapdragons and similar flowers.

THE FOLLOWING SPECIES OF ORNAMENTALS, SHRUBS AND FLOWERS CAN BE TREATED:

DECIDUOUS AND EVERGREEN TREES AND SHRUBS — Abelia, arbor vitae, Ash, Azalea, Barberry, Birch, Boxwood, Camellia, Chestnut, Cotoneaster, Cottonwood, Crab Apple, Cypress, Deutzia, Dogwood, Elaeagnus, Elm, Euonymus, Fir, Forsythia, Gum, Hawthorne, Holly, Hydrangea, Japanese andromeda, Juniper, Lantana, Lilac, Locust, Magnolia, Maple, Mock orange, Mountain Laurel, Oak, Pachistima, Pine, Pittosporum, Podocarpus, Poplar, Privet, Redbud, Rhododendron, Rose, Russian olive, Spirea, Spruce, Sycamore, Tree Peony, Tulip tree, Variegated privet, Viburnum, Walnut, Weigela, Willow, Yew.

GROUND COVERS: Ballic Ivy, Boston Ivy, English Ivy, Honeysuckle, Pachysandra, Wilson Ivy.

HERBACEOUS PLANTS: Ageratum, Alyssum, Aster, Babysbreath, Bellflower, Bleeding heart, Blood leaf, Bugloss, Candle Larkspur, Candytufts, Cinquefoil, Chrysanthemum, Columbine, Coleus, Coreopsis, Coral bells, Cosmos, Cuphea, Dahlia, Delphinium, Evening primrose, Fernleaf yarrow, Feverfew, Forget-me-not, Four-O-Clock, Foxglove, Gaillardia, Geranium, Gladiolus, Gold dust, Golden marguerite, Golden tuft, Iris, Lavendercotton, Lily, Lupine, Marigold, Morning glory, Mother-of-thyme, Mourning bride, Mourning pinks, Nasturtium, Orpine, Peonies, Petunia, Pink heath, Poker plant, Purple coneflower, Rose moss, Salvia, Scarlet sage, Snapdragon, Stone Crop, Strawflower, Sundrops, Sunflower, Sweet Pea, Verbena, Virginia Spiderwort, Wall Flowers, Wormwood, Zinnia.

Strawberries: (New Plantings): Treat after transplanting at the rate of ½ lb. per 60 square feet. (Established plantings): Treat in the fall and early spring at the rate of ½ lb. per 60 square feet. Do not apply after first bloom.

Vegetables: Apply to soil surface at 4½ lbs. product per 500 square feet (0.9 lb. per 100 square feet). Beans (field dry), broccoli, brussel sprouts, cabbage, cauliflower, collards, garlic, kale, mustard, onion, potato (whole and cut pieces), sweet potato, turnip, yam — Apply at time of seeding or immediately after transplanting. Lettuce — Apply one application only, from 1 to 3 weeks after seeding. Cantaloupe, cucumber, honeydew melon, squash, watermelon — Apply from 4 to 6 weeks after seeding. Egg plant and peppers — apply promptly after transplanting. On tomato transplants, apply 4 to 6 weeks after transplanting.

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Do not take internally. Avoid contact with eyes. Wash after handling or using. Do not contaminate feed or food-stuffs. Do not store near seeds, fertilizers or pesticides. Do not graze treated areas or feed plants parts to livestock.

CONTAINER DISPOSAL

Do not reuse container. Wrap and put in trash collection.

THE FOLLOWING PLANTS MAY BE INJURED — DO NOT USE NEAR: Bugle weed (ajuga), button pinks, carnations, dianthus, germander, geum, ice plants (mesembryanthemum), pansy, phlox, sweet william, telanthera, violets. Use around tulips, amaryllis and hosta is questionable.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

EPA Reg No 2169-180

EPA Est 2169-KS-1

Patterson GREEN-UP Company
DIV. CURRY CARTWRIGHT, INC.
1400 UNION AVE., KANSAS CITY, MO. 64101

BEST AVAILABLE COPY