

GRASSHOPPERS: Use 2½ lbs. in sufficient water to cover 1 acre (1 tablespoonful per gallon of water). The use of this product for grasshopper control should be limited to waste lands. Do not feed treated forage to animals or graze untreated areas.

SOIL INSECT CONTROL

Do not apply to soil to be used for growing root crops within 3 years after treatment.

CORN ROOT WORM AND SEED CORN MAGGOT (On Corn): Use 4 to 8 lbs. in sufficient water to cover 1 acre, a week to 10 days before planting time. Cultivate into the top 3 inches of soil. Do not apply as a row or band treatment.

WIREWORMS (On Corn): Use 4 to 8 lbs. in sufficient water to cover 1 acre, cultivating into the top 6 to 9 inches of soil. This is best done by discing, then plowing and discing again.

WARNING: Harmful if swallowed! May be absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Avoid application to feed, forage or feed or drinking troughs. Do not apply to animals.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning. (Keep out of smoke.)

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Apply this product only as specified on this label.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

PATTERSON'S

12% BHC

WETTABLE POWDER

Contains 12% Gamma BHC

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride 12.0%

Other isomers of benzene hexachloride 54.7%

INERT INGREDIENTS 33.3%

TOTAL.....100%

WARNING

Keep out of reach of children. SEE OTHER WARNINGS ON BACK PANEL.

2169-200
ACCEPTED
 DEC 4 1972
 THE FEDERAL INSECTICIDE AND FUNGICIDE BOARD
 REGISTERED UNDER THE FEDERAL INSECTICIDE ACT
 TO ATTACHED 30

1-LB. NET WEIGHT

PATTERSON CHEMICAL COMPANY
 1400 UNION AVENUE KANSAS CITY, MISSOURI

DIRECTIONS FOR USE

APPLE: APHIDS—1½-2 lbs. per 100 gallons of water (½ tablespoonful per gallon of water). Application may be repeated as necessary, but do not use within 60 days of harvest or after the fruit starts to form. Do not apply more than 30 lbs. total product per acre.

PEACH: PLUM CURCULIO—1½-2 lbs. per 100 gallons of water (½ tablespoonful per gallon of water). Apply first when about one-half of shucks have fallen, a second application when all of shucks have fallen, and a third application about 7 days later. Do not apply more than 25 lbs. total product per acre. Do not use within 60 days of harvest.

STRAWBERRIES: SPITTLE BUGS, STRAWBERRY WEEVIL—1½ lbs. per 100 gallons of water (1 teaspoonful per gallon of water). Apply only before blossoms open. Apply 100 gals. finished spray per acre.

VEGETABLE CROPS RAISED FOR SEED (Cabbage, Celery, Lettuce): Aphids—2½ lbs. per 100 gallons of water (¾ tablespoonful per gallon of water). Apply 100 gals. finished spray per acre. **NOTE:** Do not apply to celery after seedling stage. Do not apply to lettuce after thinning.

EPA Registration No. 2169-200

LABOR SAVING CHEMICALS