

CANTALOUPE, HONEYDEW MELON, WATERMELON AND SQUASH: Pickleworm, melon worm, leafhopper, aphid, squash bug, flea beetle, cucumber beetle, anthracnose, downy mildew, Alternaria blight. (Do not use closer than 7 days on Summer squash) 1 day

LETTUCE: Leafhoppers, armyworm, stinkbug, downy mildew. (Do not use on leaf lettuce within 14 days of harvest.) 7 days

PEPPERS: Tomato fruitworm, flea beetle, Fall armyworm, leafhopper, lygus bug, anthracnose 1 day

EGGPLANT: Lace bug, tomato hornworm, anthracnose, leaf spot, early and late blight 1 day

TOMATOES: Early and late blight, Septoria leaf spot, anthracnose, tomato fruitworm, flea beetle, aphid, leafhopper, tomato hornworm, aphid, Colorado potato beetle, psyllid, exposed thrip, lace bug and lygus bug 1 day

NOTICE: Buyer assumes all risk and liability of use, storage and or handling of this material not in accordance with the terms of this label

NET CONTENTS: 1 POUND

PATTERSON'S®

TOMATO DUST

INSECTICIDE FUNGICIDE

MAY BE USED UP TO DAY OF HARVEST ON TOMATOES AND MANY LISTED VEGETABLES

ACTIVE INGREDIENTS	
Zineb (zinc ethylene bisdithiocarbamate)	3.9%
Captan (N trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	2.5%
Carbaryl (1-naphthyl N-methylcarbamate)	2.0%
Rotenone	1.0%
Other cube extractives	1.6%
INERT INGREDIENTS	
	89.0%
	100.0%

CAUTION: Keep out of reach of children.

See additional cautions on side panels

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid contact with skin. Wash hands before eating or handling food. Change to clean clothing daily.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning equipment or disposal of wastes. Do not reuse container. Destroy when empty. Containers should be disposed by punching hole in them and burying with wastes or by burning (keep out of smoke).

Note for Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is an antidote.

DIRECTIONS

Apply Tomato Dust thoroughly, covering both upper side and under side of leaves and stems of plants as well as possible. Sift from this shaker-top cannister, or better still, apply with a dust gun. Apply Tomato Dust when disease or insects threaten and repeat at 7 to 10-day intervals to keep foliage covered. Dust on wet or dry foliage, but try to dust when air is still.

Tomato Dust may also be used as an aid in the control of insects and plant diseases listed below:

TIME TO STOP USE BEFORE HARVEST

BEANS: Aphid, Japanese beetle, exposed thrip, Mexican bean beetle, leafhopper, flea beetle, corn earworm, lygus and stinkbug, downy mildew, rust and anthracnose 7 days

CARROTS: Leafhopper, webworm, flea beetle, spittlebug, leaf blight, Cercospora leaf spot 7 days

CORN (SWEET): Corn earworm, fall army worm, grasshopper, leaf blight. (Do not feed forage to dairy animals or animals being finished for slaughter. Apply to corn silks every 3 days until dry for earworm control.) 7 days

CUCUMBERS: Pickleworm, leafhopper, aphid, squash bug, flea beetle, cucumber beetle, anthracnose, downy mildew, Alternaria blight 1 day

LABOR SAVING CHEMICALS

stinkbug, downy mildew. (No not use on leaf lettuce within 14 days of harvest.) 7 days

PEPPERS: Tomato fruitworm, flea beetle, Fall armyworm, leafhopper, lygus bug, anthracnose 1 day

EGGPLANT: Lace bug, tomato hornworm, anthracnose, leaf spot, early and late blight 1 day

TOMATOES: Early and late blight, Septoria leaf spot, anthracnose, tomato fruitworm, flea beetle, aphid, leafhopper, tomato hornworm, aphid, Colorado potato beetle, psyllid, exposed thrip, lace bug and lygus bug... 1 day

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

NET CONTENTS 1 POUND

3-30-72
2167-205

TOMATO DUST

INSECTICIDE FUNGICIDE

MAY BE USED UP TO DAY OF HARVEST ON TOMATOES AND MANY LISTED VEGETABLES

ACTIVE INGREDIENTS:

Zineb (zinc ethylene bisdithiocarbamate)	3.9%
Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	2.5%
Carbaryl (1-naphthyl N-methylcarbamate)	2.0%
Rotenone	1.0%
Other cube extractives	1.6%
INERT INGREDIENTS	89.0%
	100.0%

CAUTION: Keep out of reach of children.

See additional cautions on side panels.

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid contact with skin. Wash hands before eating or handling food. Change to clean clothing daily.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning equipment or disposal of wastes. Do not reuse container. Destroy when empty. Containers should be disposed by punching holes in them and burying with wastes or by burning (keep out of smoke).

Note for Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

PATTERSON CHEMICAL COMPANY

1400 UNION AVENUE

KANSAS CITY, MISSOURI

EPA Reg. No. 249-20

Dust on wet or dry foliage, but when air is still.

Tomato Dust may also be used as the control of insects and plant diseases below:

TIME
BEP

BEANS: Aphid, Japanese beetle, posed thrip, Mexican bean beetle, leafhopper, flea beetle, corn earworm, lygus and stinkbug, downy mildew, rust and anthracnose

CARROTS: Leafhopper, webworm, beetle, spittlebug, leaf blight, Cercospora leaf spot

CORN (SWEET): Corn earworm, army worm, grasshopper, leaf blight (Do not feed forage to dairy animals or animals being finished for slaughter. Apply to corn silks every 3 days until dry for earworm control.)

CUCUMBERS: Pickworm, leafhopper, aphid, squash bug, flea beetle, cucumber beetle, anthracnose, downy mildew, Alternaria blight

LABOR SAVING CHEMICALS