

Statement of Practical Treatment—In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION
Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not use when weather conditions favor drift from treated areas.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store other pesticides or acids. Do not reuse empty container. Wrap container and put in trash collection.

GENERAL INFORMATION

This product may be used for the control of broadleaved weeds in lawns, ornamental turf, vacant lots and flower rows. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65 degrees to 80 degrees F. Do not apply on a windy day. If applied during extreme temperatures weed control will be erratic. Do not apply if rain is expected in 2 hours. Application may be made with pump, knapsack or hose attached sprayers. Do not treat newly seeded lawns until after first cutting. Make sure to add Patterson's Spreader Sticker to the spray for best weed control.

Avoid spray drift to susceptible plants, as this product may injure beans, cucumbers, grapes, pears, tomatoes, and other vegetables, flower and ornamental plants. Control sprays applied at low pressure are less likely to drift. Do not exceed pre-



DANDELION

and

ROADLEAF WEED KILLER

ACTIVE INGREDIENT

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* 22.0%

INERT INGREDIENTS

Total 78.0%

*2,4-Dichlorophenoxyacetic Acid Equivalent 18% or 1.67 lbs per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautionary Statements and Statement of Practical Treatment

EPA Reg. No. 219-20 EPA P. 219-20-1

GREENUP

scribed dosage rates. Do not use equipment not for this product to apply insecticides or fungicides to susceptible plants or seeds. Do not use on grass seed crops or on turf containing bent grasses. Do not apply to Dichandra, St. Augustine, Carpet grass or clover lawns. Do not use on new grass seedlings less than two months old. Clover present in turf may be damaged. Do not contaminate water used for irrigation or domestic purposes.

NOTICE: Bayer assumes all risk and liability for use, storage, and/or handling not in accordance with the terms of this label.

LAWNS AND ORNAMENTAL TURF: For control of susceptible weeds such as Beggar-Tick, Beck-horn, Henbit, Horseweed, Annual Morning Glory, Lamb's Quarter, Mustard, Peppercorn, Plantain, Ragweed, Shepherd's Purse, and Sunflower use 2 tablespoons per gallon of water and apply either as a spot spray or over-all spray. (One gallon will usually cover about 500 sq. ft. For hard-to-kill weeds such as Spurge, Knotweed, and Chickweed repeat as necessary for control.

VACANT LOTS AND SIMILAR NON-CROP AREAS: For broadleaved weed control in these areas use 2 tablespoons per gallon of water to thoroughly wet all undesirable foliage. One gallon of spray should cover 200 to 500 sq. ft. depending on density of foliage. Make sure to use spreader sticker. Repeat each 2 to 3 weeks as necessary to control any regrowth.

SPOT SPRAYING OF WEEDS: Use a pressure sprayer or a hand gun type sprayer to treat individual weeds at anytime. New grass may be injured in these spots. Add 3 tablespoons per gallon of water making sure to use a spreader sticker. Wet weeds. Repeat as needed for control.

8 OUNCES TREATS UP TO 4,000 SQ. FT.

OVERALL SPRAYING OF WEEDS		
Amount of Spray	Amount of Water	To Cover
2 tbsp (1 oz)	1 gallon	500 Sq. Ft.
4 tbsp (2 oz)	2 gallon	1,000 Sq. Ft.
8 tbsp (4 oz)	4 gallon	2,000 Sq. Ft.
16 tbsp (8 oz)	8 gallon	4,000 Sq. Ft.

NET CONTENTS ONE PINT

35