

PREPARATION FOR USE

Mix with water in the ratio of 4 teaspoons per gallon. Repeat applications at 7 to 14 day intervals for continued control.

ORNAMENTAL INSECT CONTROL Use 4 teaspoons per gallon of water and apply when insects or their damage first appear. Spray thoroughly to cover both sides of foliage. Repeat weekly or as required to control aphid, bagworm, birch leaf miner, blister beetle, flea beetle, Japanese beetle, June beetle, lace bug, leafhopper, leaf roller, mealybug, mimosa web worm, oak leaf miner, periodical cicada, 17 year locust, plant bug, psyllid, rose aphid, rose slug, tent caterpillar, thrip, exposed, and willow leaf beetle on flowers, shrubs and trees. Do not apply to Boston Ivy.



VEGETABLE INSECT CONTROL Refer Caution No. 2. Use 4 teaspoons per gallon of water on young plants, 8 teaspoons per gallon on mature plants to control corn earworms, European corn borer, fall armyworm, fleabeetle, Japanese beetle, and leafhopper on Corn, bean leaf beetle, cucumber beetle, fleabeetle, Japanese beetle, leafhopper, velvet bean caterpillar and western bean cutworm on Beans; green lima, snap, cowpeas, and black-eyed peas; flea beetle, harlequin bug, leafhopper, armyworm, imported cabbage worm, corn earworm, tarnished plant bug and stink bug on Carrots, Chinese Cabbage, Collard, Dandelion, Endive, Garden Beets, Horse radish, Kale, Lettuce. Refer Caution No. 1. Mustard, Greens, Parsley, Parsnips, Rutabaga, Salsify, Spinach, Swiss Chard and Turnips.

Use only the low rate therefore 4 teaspoons per gallon, 2 quarts per 100 gallons water for pickleworm, melonworms, cucumber beetle, flea beetle, leafhopper and squash bug on Cucumbers, Melons, Pumpkins and Squash. See Caution No. 1.

Use 4 tsp. per gallon of water for leafhopper, Colorado Potato Beetle, flea beetle, leafhopper, European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, tarnished plant bug and stink bug on Potatoes, Tomatoes, Eggplant and Pepper; corn earworm and stink bug on Okra, fleabeetle, Harlequin bug, armyworm, imported cabbage worm, and corn earworm on Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi.

CAUTION NO. 1: To avoid possible injury to lettuce and young cress, do not apply when foliage is wet, when rain or excessive humidity is expected within the next 2 days.

NOTE: Apply this product to the trunk of trees and shrubs only when a severe pest problem exists. For control of European Corn Borer, spray thoroughly into the whorls of plants and about the fruit. Repeat at weekly intervals until the borer is controlled. For Fall Armyworm, apply to silk as soon as it appears. Repeat at 7 to 14 day intervals.



For Use on Flowers, Vegetables, Fruits, Lawns, Shrubs and Ornamentals

ACTIVE INGREDIENTS:	
Carbaryl, 1-Naphthyl N-methylcarbamate	24.4
INERT INGREDIENTS	75.6
	100.0

Contains 1 lbs. actual carbaryl per gallon.
Sevin, registered trade mark of Union Carbide Corp.
for carbaryl insecticide U.S. Pat. Nos. 2,503,478, 3,009,855

CAUTION: Keep out of reach of children.

See other Caution on Side Panel

E.P.A. Reg. No. 2169-201-1000

CAUTION NO. 2: Apply at 7 to 10 day intervals to all above ground parts, including sprouts and flowers, radish, turnips, rutabaga, endive, green tops, kale, leaf lettuce.

FRUIT TREES: Apples. Apply to non-bearing week intervals. On bearing trees, apply at 10 to 14 day intervals when used at 10 to 15 days after harvest. Apply on the above schedule and observe lesser peach tree borer and June beetles. Do not apply to peaches.


FRUIT REDUCTION: On desired on Crabapple. 6 teaspoons per gallon 10 to 15 days after full bloom. Do not expect more than 50% reduction.

LAWN INSECT CONTROL: Use 15 gallons of water per 100 sq. ft. for lawn attached to pre-bud ants, carwigs, fire ants, brown dog tick, and other lawn pests. Repeat watering every 10 to 15 days.

CAUTION: Handle with care. Do not take internally. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly with soap and water after use. Do not eat, drink, or smoke while using this product.

Do not use on lawns where children play. Do not use on lawns where pets are kept. Do not use on lawns where livestock are kept.

NOTE: For control of European Corn Borer, spray thoroughly into the whorls of plants and about the fruit. Repeat at weekly intervals until the borer is controlled. For Fall Armyworm, apply to silk as soon as it appears. Repeat at 7 to 14 day intervals.



For Use on Flowers, Vegetables, Fruits, Lawns,
Shrubs and Ornamentals.

ACTIVE INGREDIENTS:

Carbaryl, 1-naphthyl N-methylcarbamate 24.4%

INERT INGREDIENTS 75.6%

100.0%

Contains 2 lbs. actual carbaryl per gallon.
Sevin, registered trade-mark of Union Carbide Corp.
for carbaryl insecticide U.S. Pat. Nos. 2,903,478; 3,009,855

CAUTION: Keep out of reach of children.

See other Cautions on Side Panel

E.P.A. Reg. No. 2169-207-1-72

CAUTION NO. 2: Apply when insects or the insects appear. Repeat at 7 to 10 day intervals or as necessary. Do not apply within 3 days of harvest on all above crops, except 3 days for cabbage, broccoli, Brussels Sprouts, cauliflower, kohlrabi, head lettuce, garden beet, root, parsnips, radish, parsnips, rutabaga, salsify and turnip, and 14 days for collards, dandelion, endive, garden beet, tops, spinach, Swiss Chard, and turnip tops, kale, leaf lettuce, mustard, green parsley and salsify tops.

FRUIT TREES: Apples and Peaches: Use 4 teaspoons per gallon of water. Apply to non-bearing trees 7 to 10 days after full leaf and repeat at 2-3 week intervals. On bearing trees, to avoid thinning of fruit delay use at least 30 days after bloom. Use Patterson's Fruit Tree Spray earlier. Controls: when used at 10 to 14 day schedule: green apple aphid, fruit tree and red banded leaf roller, Bagworms, plum curculio, and codling moth. Do not apply on the above mentioned fruit trees later than 1 day before fruit harvest.

Spray Peaches, and Plums at same rate: 4 teaspoons per gallon, same schedule and observe precautions for bearing trees. Controls: plum curculio, lesser peach tree borer, Oriental fruit moth, twig borer, fruit tree roller, and June beetles. Do not apply later than 1 day before fruit harvest.

FRUIT REDUCTION ON ORNAMENTAL TREES: When fruit production is not desired on Crabapple, Plum, Mulberry, and other decorative trees, spray 8 teaspoonfuls per gallon, particularly to the blooms, between 10 and 15 days after full bloom. One treatment is prescribed and complete fruit drop should not be expected.

LAWN INSECT CONTROL: Use 1 pint per gallon or 16 tablespoonfuls per 16 to 15 gallons of water and apply evenly to each 500 square foot. Use hose attached or pressure type sprayer to control: fire, cutworm, chinch bug, ants, earwigs, fall armyworm, fleas, leaf hopper, millipede, mosquito, and brown dog tick. For best results, mow lawn first and apply after rain or watering. Repeat 2 or 3 weeks later, if necessary. For Sod web worms apply May 15th and repeat at 21 day intervals through August.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing fumes. Do not take internally. Avoid contact with skin. Wear long sleeved clothing and clean after spraying. Wash thoroughly after handling. Wash hands and face before eating. To avoid injury to trees, foliage, do not spray when foliage is wet or when rain or excessive humidity is expected. The product is highly toxic to bees, wasps, etc. Do not treat flowers, shrubs, or crops. Protective information may be obtained from Cooperative Agricultural Extension. Do not contaminate food, feed, water, or other articles. Destroy contents by pouring contents into water and flushing out of tank.

NOTE FOR PHYSICIANS: Carbaryl is a carbamate. It is a central nervous system inhibitor. Atropine is antidote.

NOTICE: Before using, check the label for directions and cautions. Do not use if label is missing or illegible.

CONTENTS: 1 QUART