

CAUTION NO. 2: Apply when insects or their damage appear. Repeat at 7 to 10 day intervals, or as necessary. Do not apply within 1 day of harvest on all above crops, except 3 days for cabbage, broccoli, brussels sprouts, cauliflower, kohlrabi, head lettuce, garden beet (root), horseradish, radish, parsnips, rutabaga, salsify and turnip, and 14 days for collards, dandelion, endive, garden beet (tops), spinach, swiss chard, and turnip (tops); kale, leaf lettuce, mustard greens, parsley and salsify tops.

FRUIT TREES: Apples and Pears. Use 4 teaspoons per gallon of water. Apply to non-bearing trees 7 to 10 days after full leaf and repeat at 2-3 week intervals. On bearing trees, to avoid thinning of fruit delay use at least 30 days after bloom. Controls, when used at 10 to 14 day schedule, green apple aphid, fruit tree and red banded leaf roller, Bagworms, plum curculio, and codling moth. Do not apply on the above mentioned fruit trees later than 1 day before fruit harvest.

Spray Peaches, and Plums at same rate (4 teaspoons per gallon), same schedule and observe precautions for bearing trees. Controls plum curculio, lesser peach tree borer, Oriental fruit moth, twig borer, fruit tree rollers and June beetles. Do not apply later than 1 day before fruit harvest.

LAWN INSECT CONTROL: Use 1/2 pint (8 ounces or 16 tablespoonfuls) in 10 gallons of water and apply evenly for each 500 square feet. Use backpack or pressure-type sprayer to control sod webworm, chinch bug, mole cricket, fall armyworm, flea, leafhopper, millipede, mosquito, and other lawn insects. For best results, mow lawn first and apply after rain or irrigation. For sod webworms apply through August.

SEVIN*

LIQUID CONCENTRATE

For Use on Flowers,
Vegetables, Fruits, Lawns,
Shrubs and Ornamentals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Mix with water as directed. Keep spray mixture well agitated. Repeat applications at 7 to 14 day intervals for continued control.

ORNAMENTAL INSECT CONTROL. Use 4 teaspoonfuls per gallon of water (equals 2 quarts per 100 gallons) and apply when insects or their damage first appear. Spray thoroughly to cover both sides of foliage. Repeat weekly or as required to control Aphids, Bagworms, Birch Leaf Miner, Blister Beetle, Flea Beetle, Gypsy Moth Larvae, Japanese Beetle, June Beetle, Lace Bugs, Leafhoppers, Leaf Roller, Mealybugs, Mimosa Webworm, Oak Leaf Miner, Periodical Cicada (17 year Locust), Plant Bugs, Psyllid, Rose Aphid, Tent Caterpillar, Thrip (exposed stages), and Willow Leaf Beetle on Flowers, Shrubs, Ornamental and Forest Trees. Do not apply to Boston Ivy.

VEGETABLE INSECT CONTROL: (Refer Caution No. 2) Use 4 teaspoons per gallon of water on young plants, 8 teaspoons per gallon on mature plants to control corn earworms, European corn-borer, fall armyworm, flea beetle, Japanese beetle, and leafhopper on Corn; bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhopper, velvet bean caterpillar and western bean cutworm on Beans; (green, lima, snap, cowpeas, and blackeye peas); flea beetle, harlequin bug, leafhopper, armyworm, imported cabbage worm, corn earworm, tarnished plant bug and stink bug on Carrots; Chinese Cabbage Collard, Dandelion, Endive, Garden Beetle, Horseradish, Kale, Lettuce (Refer Caution No. 2); Mustard Greens, Parsley, Parsnips, Rutabagas, Salsify, Spinach, Swiss Chard and Turnip.

ACCEPTED
 APR 12 1984
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under EPA Reg. No. 2169-207

2169-207

49612179

111