

Feb-16-1973

GENERAL INFORMATION

Patterson's Dandelion Killer may be used for control of broad-leaved weeds in lawns, turf, vacant lots, and fence rows. It is most effective when weeds are small and actively growing and should be applied when the temperatures range from 65° to 90° F. Do not apply on a windy day. If applied during extreme temperatures weed control will be erratic. Do not apply if rain is expected within two hours. Application may be made with either power, knapsack or hose-attached sprayers. Do not treat newly seeded lawns, until after first cutting. Make sure to add Patterson's Spreader Sticker to the spray for best weed control.

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TREES: For control of susceptible weeds, as named on the front panel use 2 tablespoon per gallon of water and apply either as a spot or over-all spray. (One gallon will usually cover about 500 sq. ft.) For hard-to-kill weeds such as Spurge, Knotweed, and Chickweed repeat as necessary for control, or use Patterson's Chickweed Killer containing Silvex.

VACANT LOTS AND WASTE AREAS: For broadleaved weed control in these areas, use 3 tablespoonfuls per gallon of water to thoroughly wet all undesirable foliage. One gallon of spray should cover from 200 to 500 sq. ft. depending on denseness of foliage. Make sure to use spreader sticker. Repeat each 2-3 weeks as necessary to control any regrowth.

SPOT SPRAYING OF WEEDS: Use a pressure sprayer or a hand gun type sprayer to treat individual weeds at anytime. New grass may be injured in these spots. Add 3 tablespoons per gallon of water making sure to use a spreader sticker. Wet weeds. Repeat as needed for control.

NET CONTENTS—1 PINT

PATTERSON'S

DANDELION KILLER

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid²

22.0%

INERT INGREDIENTS

78.0%

Total 100.0%

²2,4-Dichlorophenoxyacetic Acid Equivalent 18.3%

CAUTION: Keep out of reach of children.

See other cautions on side panel.

PATTERSON CHEMICAL COMPANY

1900 UNION AVENUE

KANSAS CITY, MISSOURI

EPA Reg. No. 2167-206

OVERALL SPRAYING OF WEEDS

Amount of Spray	Amount of Water	To Cover
2 tbsp. (1 oz.)	1 Gallon	500 sq. ft.
4 tbsp. (2 oz.)	2 Gallon	1000 sq. ft.
8 tbsp. (4 oz.)	4 Gallon	2000 sq. ft.
16 tbsp. (8 oz.)	8 Gallon	4000 sq. ft.

Use a pressure sprayer or Patterson Hose-End Sprayer. Add amount of spray to sprayer bottle, fill to desired level and spray. Make sure of square footage before spraying. Treat on a calm day.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, or clothing. In case of contact, wash off promptly with plenty of water.

Avoid spray drift to susceptible plants, as this product may injure, beans, cucumber, grape, rose, tomato, and other vegetable, flower and ornamental plants. Coarse sprays applied at low pressure are less likely to drift. Do not exceed prescribed dosage rates. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not use equipment used for applying this product to apply insecticides or fungicides to susceptible plants or seeds. Do not use on grass seed crops or on turf containing bent grasses. Do not apply to Dichondra, St. Augustine carpet grass or clover lawns. Do not use on new grass seedings less than two months old. Clover present in turf may be damaged.

Do not contaminate lakes, streams, ponds, or water used for irrigation or domestic purposes. Do not reuse empty container. Dispose in a safe place.

NOTICE: Buyer assumes all risk and liability of use, storage, and/or handling not in accordance with the terms of this label.

Also kills other common lawn weeds, such as Beggartick, Buckhorn, Heal All, Horseweed, Annual Morning Glory, Lambs Quarter, Mustard, Peppergrass, Plantain, Ragweed, Shepherd's Purse, Sunflower

1-72 8 OUNCES TREATS UP TO 4,000 SQ. FT.

LABOR SAVING CHEMICALS