

Time to Stop Max. Dosage
Use Before Product
Harvest Per Acre

STRAWBERRIES: Meadow spittle bug,
strawberry leaf roller and straw-
berry weevil 1 Day 40 lbs.
($\frac{2}{3}$ lb./1,000 sq. ft.)

GRASSHOPPERS ON THE ABOVE CROPS: Apply at the dosages and limitations given for each crop and repeat as necessary to control grasshoppers.

ORNAMENTALS: PATTERSON'S 5% SEVIN DUST can be used on a wide range of ornamental plants, flowers, shrubs and small trees, namely, rose, carnation, gladiolus, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines. Do not use on Boston Ivy.

Apply to upper and lower leaf surface of all foliage for complete coverage to control pests as apple aphid, bagworm, birch leaf miner, blister beetle, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetle, gypsy moth, Japanese beetle, lace bugs, leaf hopper, mealybugs, rose aphid, thrips and rose slug.

CATS AND DOGS: For ticks and fleas, dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the hair. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens or puppies less than 4 months of age.

In and around dog and cat quarters: For ticks and fleas, dust liberally in and around sleeping quarters, kennels, and dog houses by means of a shaker can, hand or dust gun.

NOTE: SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment. Do not use on Boston Ivy.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid breathing of dust. SKIN CONTACT MAY BE HARMFUL. Avoid contact. Wash hands before eating. Take shower or bath after work. WEAR REGULAR OR LONG-SLEEVED WORK CLOTHING. Change to clean clothing daily. Do not apply to open blossoms. This product is toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Keep out of reach of animals. Note for Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

PATTERSON'S®

5% SEVIN* DUST

For Control of Certain Insects on Vegetables,
Strawberries, Ornamentals, and pets.

ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%
100%

*Registered trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

CAUTION:

Keep out of reach of children.
See other warnings on back panel.

NET WEIGHT 1 POUND

MANUFACTURED BY, OR TO THE SPECIFICATIONS OF
PATTERSON CHEMICAL COMPANY
1400 UNION AVENUE • KANSAS CITY, MISSOURI

E.P.A. Reg. 2169-214 3-72

DIRECTIONS FOR USE

Apply PATTERSON'S 5% SEVIN DUST when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surface of all foliage for complete coverage. For best results apply with a dust gun. Repeat as necessary, generally at 7-day intervals. Do not apply in excess. Make applications in the morning or evenings when it is cool and the foliage is dry.

FOR CONTROL OF

Time to Stop Max. Dosage
Use Before Product
Harvest Per Acre

BEANS (green beans, lima beans, snap beans, cowpeas, blackeyed): Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetles, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworms, corn earworms, stink bugs and tarnished plant bugs..... No Limit 40 lbs.
($\frac{2}{3}$ lb./1,000 sq. ft.)

CUCUMBER, MELONS, PUMPKIN, SQUASH: Pickleworm, melonworm, cucumber beetle, flea beetle, leafhopper, and squash bug. Avoid excessive applications. Do not apply to wet foliage. No Limit 20 lbs.
($\frac{1}{3}$ lb./1,000 sq. ft.)

POTATO, TOMATO, EGGPLANT, PEPPER: Colorado potato beetle, flea beetle, leafhopper, European corn borer, fall armyworm, lace bugs, tomato fruit worm, tomato hornworm, tarnished plant bugs, stink bugs..... No Limit 40 lbs.
($\frac{2}{3}$ lb./1,000 sq. ft.)

ASPARAGUS: Asparagus beetle and Apache cicada (Do not repeat application within 3 days) 1 Day 30 lbs.
($\frac{2}{3}$ lb./1,000 sq. ft.)

CARROTS (No time limit), HEAD LETTUCE, HORSE RADISH, PARSNIPS, RADISHES, RUTABAGAS: Flea beetles, harlequin bug, leafhoppers, armyworms, cabbage caterpillars, corn earworms, spittlebugs, tarnished plant bugs and stink bug 3 Days 40 lbs.
($\frac{2}{3}$ lb./1,000 sq. ft.)

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea beetle, harlequin bug, armyworms, imported cabbage worm, corn earworm 3 days 40 lbs.
($\frac{2}{3}$ lb./1,000 sq. ft.)

LABOR SAVING CHEMICALS

2169-214


3-31-1972

gpc

1/2

DATA

3-23-72



2 169-214

2/2