

## DIRECTIONS FOR USE

**CRABGRASS CONTROL:** To establish good crabgrass control in permanent turf areas, this entire bag content should be applied evenly over 7,500 square feet in northern cool grass areas and 5,000 square feet on southern warm climate grasses. Application must be made in the spring before crabgrass germinates. Patterson's Pro-Teck must be applied evenly over the turf area without skips or misses. For best results, one half of the required total amount should be spread in one direction and the remaining one half spread over the same area but at right angles to the first application.

**ANNUAL BLUEGRASS (Poa Annua) CONTROL:** Northern Areas—For cool season grasses apply the entire bag content to 7,500 square feet (6x25 lb. bag per acre). This should be done in late summer or early fall before annual bluegrass germinates. Application method is the same as described in this label under Crabgrass Control. This provides 2 pounds of actual BENEFIN per acre.

**Southern Areas—**For warm season grasses such as bermudagrass, St. Augustinegrass, and bahiagrass apply the entire content of this bag to 5,000 square feet. This should be done in late summer or early fall before annual bluegrass germinates. Application method is the same as described in this label under Crabgrass Control. This provides 3 pounds actual BENEFIN per acre.

## APPLICATION EQUIPMENT


Equipment designed to apply granular herbicides or insecticides should be used. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

	Southern	Northern
Spreader Setting: Scott (drop)	7	6
Central	9	8
Greenfield	9	8
Cyclone	5½	4½
Sears	7	6
Jackson	7	6
Agrico	7	6

Both drop-type and rotary-type spreaders may be used to apply Patterson's Pro Teck. A more uniform application can be made with spreaders which avoid concentration of the material in narrow bands and which deliver granules to an area as wide as the hopper width.

E.P.A. Est. 2169-Ks.-1

PATTERSON'S



**PRO-TECK**  
HERBICIDE

A herbicide for professional use on lawns and golf courses. When used as directed, will control crabgrass, annual blue grass (poa annua), and other grasses in established:

BLUEGRASS (Perennial)	BERMUDA GRASS
RYEGRASS (Perennial)	CENTPEDE GRASS
FESCUE	ST. AUGUSTINE GRASS
BAHIA GRASS	ZOYSIA GRASS

### ACTIVE INGREDIENTS:

N-Butyl-N-Ethyl-a-a-a-Trifluoro-2,6-

Dinitro-p-Toluidine 1.3%

INERT INGREDIENTS: 98.7%

(Relative to herbicidal content) 100.0%

BALANCE

The Trademark for BENEFIN Elanco Products Company  
a division of Eli Lilly & Company

Treats 7,500 square feet when treating for crabgrass control.

## CAUTION:

Keep out of reach of children.

See back panel for additional cautions.

**NET WEIGHT 25 LBS.**

EPA Reg. No. 2169-222

## PRECAUTIONS

Apply Patterson's Pro Teck Granular to the turf prior to the germination of weed grasses, since the material will not control established weeds. Patterson Pro Teck may thin fine-leaved fescues at rates above 112.5 pounds per acre (1.5 pounds active ingredient). Patterson Pro Teck Granular should not be applied in the spring to bluegrass planted the previous fall. Do not apply to bentgrass putting greens, or dichondra turf. Do not apply to newly sprigged areas of bermudagrass, St. Augustinegrass, or centipedegrass until these turfs are well established.

## CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of contact, flush with water. Eye irritant. Wash thoroughly with soap and water after use. Do not contaminate foodstuffs or feeds.

Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock. Destroy bag when empty. Do not reuse. Dispose in a safe place.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate any body of water by direct applications, cleaning of equipment, or disposal of wastes.

## GENERAL DIRECTIONS

Patterson's Pro Teck is a pre-emergence herbicide which is applied directly to established turf for the control of smooth crabgrass, hairy crabgrass, goosegrass (silver crabgrass), watergrass (barnyard grass), yellow and green foxtail, and annual blue grass (poa annua.)

Patterson's Pro Teck controls annual grass weeds by killing their seeds as they germinate. It does not control established weeds. When it is applied as directed the following established grasses are tolerant to Patterson's Pro Teck: perennial bluegrasses, perennial ryegrass, centipedegrass, fescue, zoysiagrass, bermudagrass, St. Augustingrass, and bahiagrass.

Apply Patterson's Pro Teck to turf before germination of annual grass weeds. Application time will vary by area, depending on the weed grasses to be controlled and their time of germination. Do not reseed within 42 days after application.

**NOTICE:** Buyer assume all risk and liability of use, storage, and/or handling of this material not in accordance with the terms of this label.