BEST DOCUMENT AVAILABLE



COMMENTS

EPA Letter Dated:

JUN2 2 1984

Under the Federal Insecticide, Pungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. Ha 2107 - 232

WETTABLE
DUSTING SULFUR

Use As A Dust or Liquid Speny

FUNGICIDE - MITICIDE

	Sulfur
	INERT INGREDIENTS: 8.0%
	Total 100.0%
	KEEP OUT OF REACH OF CHILDREN 10 PT
	CAUTION
	See Side Fanel For Additional Precautionary Statements
EPA	Reg. No. 2169-232 EFA But. No. 2169-KD-1

Net Content:

ACTIVE INGREDIENT:

FAITERION GREEN-UP COMPANY Farry Continues F. x 0015 Karate City, Mo. - 7.101

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

Wettable Dusting Salfar MPA Rev. No. 1100 232 . Page 2 of 3

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTE - Applications of this material should not be made prior to or during periods of excessively high temperatures or within 2 weeks of an oil spray. On citrus do not make an oil application within 60 days prior to or after a sulfur application.

HOW TO MIX - With spray tank half full and the agitator running sift in the required amount of Sulfur. Continue filling and keep agitator running. If applied with a pressure hand sprayer agitate solution frequently during application. This material is guaranteed to pass 95% thru a 325 mesh screen. Use 4/tablespoonfuls per gallon of water for all spray instructions.

FOR DISEASES & MITES

FOR USE AS A SPRAY, For control of BROWN ROT AND PEACH SCAB on Peaches; Use 4 level tablespoonfuls per gallon of water. (equal to 6 1/4 lb. per 100 gallons of water.) Apply to trees in the pre-bloom, bloom, and post bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. For BLACK SPOT on Roses and LEAF SPOT on Phlox and Hollyhocks. Treat when first signs of disease appears and repeat at 7 to 10 day intervals. For RUST (Puccinia & Uromyces) on Crysanthemums, Carnations, Snapdragons, and Asparagus apply at first signs of disease and repeat as needed. For POWDERY MILDEW on Roses, Lilacs, Chrysanthemums, Dahlias, Calendulas, Delphinium, Phlox, Sweet Peas, Peas, Beans, Currants, Gooseberries, and Peaches; apply at first signs of disease and repeat in 7 to 10 days. For control of TWO SPOTTED MITES & EUROPEAN RED MITE on Almonds, Avacados, Beans, Evergreens (Junipers), Gooseberries, Figs, Peas, Plums, Prunes, and Citrus Fruits; apply in early summer when mites tend to build up and repeat 's needed for control or at 7 to 10 day intervals. See PE-ENTRY STATEMENT. or workers under CAUTION.

FOR USE AS A DUST - This product may be used as a dust directly to the above mentioned plants for the control of those respective diseases & mites. Dust using approximately 1 pound of product per 1,000 sq.ft. of plants; (equal to approximately 44 lbs. per acre.) This material may be applied directly from the shaker top canister or with a mechanical or power duster. Apply to deposit a very fine layer of sultar dust to all parts of the plant. Do not over cose since foliage burn may result. Sor NOTE above pertaining to oil sprays.

PREVENTION AND CONTROL OF CHIGGERS - To prevent chigger bites dust this material directly on lower legs and ankles and the bottoms of clothing likely to be exposed to vegetation carrying chiggers. After exposure wash legs and ankles and exposed clothing with soap and water. TREATMENT OF VEGETATION-For Chigger control apply 1 pound of this product to treat 1,000 to 1300 Square Feet of vegetation. Apply to lawns, flower gardens, shrubs, and any place where chiggers may be hiding or where people contact foliage. Make a repeat application in about two weeks for continued control.

BEST DOCUMENT AVAILABLE

Page 3 of 3

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS

CAUTION - Harmful if swallowed. Avoid breathing dust or spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes or skin. Do not contaminate feed or foodstuffs. Do not apply to any area not specified on this label. RE-ENTRY STATEMENT - Do not re-enter treated areas within 24 hours after application unless appropriate protective clothing is worn.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

Do not apply when weather conditions favor drift from the areas treated. Do not apply directly to water or wetlands.

CAUTION- Some fruits and plants are susceptible to injury from Sulfur under certain climatic conditions. The user is advised not to use Sulfur on any crop or plant unless local use has proved that sulfur does not damage plants in that locality. Sulfur may burn plant foliage when the temperature is 90°F or higher in conjunction with high humidity. Do not apply at such times. Do not make application within two weeks before or after an oil spray. On Citrus do not apply, within 60 days prior to or after an oil spray.

NOTICE - Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

多少少

JUN 22 1984

Patterson Green-Up Company 1331 Union Ave. Kansas City, HS 64101

Attention: H. R. McLane

Gentlemen:

- 🚅

Subject: Green-Up Wettable Dusting Sulfur EPA Registration No. 2169-232 Your Letter Dated Pebruary 23, 1984

The labeling referred to above, submitted in connection with registration under the Pederal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Pive copies of the finished labeling must be submitted.

For asparagus, indicate that application should not be made until after cutting stops.

A stamped copy is enclosed for your records.

Sincerely yours,

1

Henry M. Jacoby Product Manager (21) Fungicide-Herbicide Branch Registration Division (TS-767)

Enclosure

RD/FHB:DCR-03272:Jacoby:eg:Raven:557-2226:RD-50:4/20/84:Del.5/4/84

BEST DOCUMENT AVAILABLE

CONCURRENCES										
-										
FURNAMED										
DATE										
MA Form 1320 1 4 81 OFFICIAL FILE COP										

• • • • • •

• •