

NUISANCE PESTS IN OUTSIDE AREAS: For control of flies, ants, cockroaches, boxelder bugs, clover mites, spiders and earwigs mix 8 ounces in 1½ gallons water and thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling, foundations, walks, garbage cans, and into cracks and other places where insects may hide. Apply when insects appear and repeat treatment when needed.

For control of ticks and chiggers spray weeds and brushy incrop areas around home. Also treat lawn for chiggers.

NOTE: Do not use on certain ferns, poinsettia, hibiscus, papaya, piles and gardenia, and orchids since injury may occur.

CAUTION: Harmful if swallowed. May be absorbed through the skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not permit children or pets to go onto sprayed grass until spray has completely dried.

FISH AND WILDLIFE CAUTION: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Do not use, pour, spill, or store near heat or open flame.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling not in accordance with the terms of this label.

**Equivalent to 0.20% of (butylcarbityl) 6-propyl-piperonyl ether and 0.05% related compounds.

*Drazinon® trademark of CIBA-GIEGY Corp.
U. S. Patent No. 2,754-243



Patterson

E.P.A. Reg. No. 2169-237 2-74

Keep Out of Reach of Children
See Side Panel for Additional Cautions

CAUTION:

ACTIVE INGREDIENTS:	
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorohidate*	12.50%
Pyrethrins	.03%
Technical Piperonyl Butoxide**	.25%
Solvent Aromatic Petroleum Distillate	79.22%
INERT INGREDIENTS	8.00%
Total	100.00%



DIRECTIONS FOR USE

FOR LAWNS: To control chinch bugs, sod webworms, bermuda mites, ants, ticks, earwigs, lawn billbugs, in bluegrass, fescue, zoysia and bermuda use 2 tablespoons (1 fluid ounce) in 1½ gallons of water to cover 65 sq. ft. One pint covers 1000 sq. ft. For millipedes double this dosage rate. Mow the lawn and thoroughly water before treatment. For billbugs water the lawn after spraying.

FLOWERING PLANTS: For roses, azaleas, mums and gladiolus to control aphids, European red mite, leafhoppers and thrips use 2 tablespoons per 1½ gallon of water. Add Patterson's Spreader Sticker to the spray and thoroughly treat foliage and flowers.

ORNAMENTAL TREES AND SHRUBS: On oaks, maple, locust, holly, boxwood, pine, spruce, and willow to control aphids, mites, scale crawlers, dipterous, leaf miners (exposed stages and bagworms) use 2 tablespoons per 1½ gallon water and thoroughly spray.

FRUIT: Use on **APPLES** for codling moth, mites, aphids and fruit tree leaf-rollers. On **PEACHES** for twig borers, scale crawlers and mites; on **STRAWBERRIES** for aphids, mites and strawberry leafrollers; on **GRAPES** for berry moth, Pacific spider mites and leafhoppers. Use 2 tablespoons in 1½ gallon water and thoroughly spray. Apply when bloom is completed and then repeat at 10 to 14 day intervals through the season but not later than 14 days on apples, 20 days on peaches, 5 days on strawberries, 18 days on grapes, before harvest or picking.

VEGETABLES: Use on tomatoes, beans, cucumbers, onions, peppers, cabbage, for control of bean beetles, aphids, imported cabbageworms, cucumber beetles and red spiders. Mix 2 tablespoons per 1½ gallon water to cover 500 sq. ft. of garden area. Apply when insect damage is seen and repeat generally at 7 to 10 day intervals. Do not apply on crops within the following days of harvest: Beans, 7 days (wait 4 days if forage is to be cut for hay). Cucumber and onions, 10 days. Peppers, 5 days. Cabbage, 7 days. Tomatoes, 1 day.