

[2169-240]

HOW TO MIX AND APPLY

Caterpillar Spray is easy to mix and apply through pressurized or hose-end sprayers. Place one-half the required amount of water in jar or tank. Add recommended amount of Caterpillar Spray while shaking or stirring. Then add balance of water and shake. A spreader sticker may be added as an aid in obtaining thorough coverage. Do not store diluted spray more than 12 hours.

Caterpillar Spray works as a stomach poison. Caterpillars must eat a small amount of treated leaf to get a lethal dose. Thorough and uniform coverage of both sides of all foliage subject to attack is essential for prompt and complete control. Within a few hours after taking a bite of treated leaf, caterpillars stop feeding. Death follows in a few days.

Apply when caterpillars (or their damage) first appear. Re-treatment at 7 to 10 day intervals may be required, especially when egg hatch occurs over an extended period of time. Use of the higher range of recommended rates provides a longer period of control.

ACTIVE INGREDIENT:

Bacillus thuringiensis, Berliner, 4,320 International Units of potency per milligram.* At least 6.75 billion viable spores per gram..... 0.86%


INERT INGREDIENTS:99.14%

*Equivalent to 1.96 billion International Units of potency per pound of this product.

®Registered Trademark, Abbott Laboratories

NOTICE: Buyer assumes all risk and liability for use, storage, and/or handling of this product not in accordance with the terms of this label.

NET WEIGHT: 4 OZ.



hose-end sprayers.
omended amount
water and shake.
coverage. Do not

a small amount
of both sides of
control. Within a
ng. Death follows

at 7 to 10 day
extended period
longer period of

ency
..... 0.86%
..... 99.14%
of this product.

d/or handling of



Ecologically Compatible
CAUTION: Keep Out of Reach of Children.
Do Not Re-Use Container. Destroy When Empty.
Distributed by:

Patterson

A Home & Garden formulation for control of leaf-eating caterpillars on shade and ornamental trees.

AMOUNT OF CATERPILLAR SPRAY TO APPLY
Ornamental and Shade trees (up to 25 feet tall)
fruit

Pest
Gypsy Moth Larvae*
Elm Spanworm
Spring and Fall Cankerworms (inchworms)
Bagworm
Fall Webworm
Red-Humped Caterpillar (California only)
*For best control of Gypsy Moth, apply when caterpillars are 1st instars). Repeat treatment if necessary.

VEGETABLES**

Crop	Pests
Broccoli, Cabbage, Cauliflower	Cabbage Looper
Beans, Collards, Cucumbers (Southern U.S.), Kale, Lettuce, Melons, Potatoes, Mustard Greens, Spinach, Turnip Greens	Imported Cabbage Worm
Celery	Cabbage Looper, Celery Looper
Grape	Grape Leaf Folder
Tomatoes	Cabbage Looper
Tobacco	Tomato Hornworm
	Budworm, Cabbage Looper
	Hornworm

**Caterpillar Spray may be applied up to the day of harvest.

E.P.A. Reg. No. 2169-240 3-73

E.P.A. Est. No. 2169-KS1