

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

Determine the amount of Broadleaf Weed Killer to be used in the sprayer by multiplying the number of gallons to be prepared by the quantity of Broadleaf Weed Killer recommended per gallon. Place this amount in the sprayer and add the required amount of water. Uniform mixing and application is essential for effective results. Weeds are most effectively controlled when in an active stage of growth—spring, summer or fall.

WEEDS IN VACANT LOTS & FENCE ROWS: For best results on thistles, ragweed, morning glory, dock and many other weeds growing in non-turf areas, spray when these weeds are in bud to flower stage. Wet all weeds thoroughly. Repeat applications may be needed for maximum control. Mix 1/2 pint Broadleaf Weed Killer in 6 gallons of water.

WOODY GROWTH IN FENCE ROWS: To control poison ivy, poison oak, poison sumac, horse nettles and brambles, dilute 5 tablespoons per gallon of water and apply as directed. Foliage spray in early summer when leaves are fully developed. Repeat as required for control.

STUMP TREATMENT: To prevent resprouting, thoroughly wet cut surfaces and all exposed surface bark with a solution of 8 oz. of Broadleaf Weed Killer, mixed with one quart of diesel oil or kerosene.



KILLS: DANDELION, PLANTAIN, DOCK, C. WEED, RAGWEED, AND MANY OTHER COMMON BROAD-LEAVED WEEDS

CAUTION

Avoid contact with skin, eyes, and clothing. Avoid mist. Do not reuse container. Destroy when empty.

AVOID INHALING SPRAY MIST

Under no circumstances should this herbicide or 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, grapes, garden crops, ornamental other susceptible crops are grown. Extreme care be exercised to prevent drifting of this material. Apply on a windy day. (Coarse sprays are less likely to drift.) Vapors from this product at very high temperatures may cause injury to susceptible plants nearby. Do not use spray equipment which has contained this product for other purposes. Do not store fertilizers, seeds, insecticides, fungicides or food. Avoid contamination of water supplies that may be used for spray, irrigation, or water susceptible crops or for domestic purposes. Do not allow animals to graze on treated turf.

Do not use, pour, spill, or store near heat or open flames. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not apply to forest lands or management areas around homes or gardens, ditchbanks, water, right-of-way, pastures or recreational areas.

NOTICE: Buyer assumes all risk and liability of storage and/or handling of this material not in accordance with the terms of this label.

12-7
E.P.A. Reg No 2160-241 E.P.A. Est No 2160

5 Teaspoons = 3 Tablespoons = 1 oz.
Net Contents 1 Quart

ACTIVE INGREDIENTS:	
Isooctyl ester of 2,4-dichlorophenoxyacetic acid*	9.4%
Isooctyl ester of Silvex [2-(2,4,5-trichlorophenoxy) propionic acid]**	4.4%
Total	13.8%
INERT INGREDIENTS	
Total	86.2%
100.0%	

* 2,4-dichlorophenoxyacetic acid equivalent—6.23% or .47 lbs. equivalent to 3.11% Silvex 2-(2,4,5-trichlorophenoxy) propionic acid for 23 lbs.

CAUTION: Keep Out of Reach of Children.
SEE OTHER CAUTIONS ON SIDE PANEL.

GREEN UP

33

ACCEPTED
20 FEB 1980
7169-241

**UTION AND
AREFULLY**

Broadleaf Weed
by multiplying
prepared by the
Killer recom-
amount in the
amount of water.
on is essential
re most effec-
active stage of

FENCE ROWS:
weed, morning
eds growing in
weeds are in
eds thoroughly.
eeded for maxi-
broadleaf Weed

ROWS: To con-
poison sumac,
5 tablespoon
as drench
when leaves
red for control.
event resprout-
es and all ex-
tion of oz. of
with on



ACTIVE INGREDIENTS:	
Isocetyl ester of 2,4-dichlorophenoxyacetic acid*	9.4%
Isocetyl ester of Silvex [2-(2,4,5-trichlorophenoxy) propionic acid]**	4.4%
INERT INGREDIENTS	
Total	86.2%
Total 100.0%	
* 2,4-dichlorophenoxyacetic acid equivalent—4.23% or .47 lbs. equivalent to 3.11% Silvex 2-(2,4,5-trichlorophenoxy) propionic acid or .23 lbs.	

CAUTION: Keep Out of Reach of Children.
SEE OTHER CAUTIONS ON SIDE PANEL

GREEN UP

33

ACCEPTED
80 FEB 1980
7169-241

KILLS: DANDELION, PLANTAIN, DOCK, CHICK-WEED, RAGWEED, AND MANY OTHER COMMON BROAD-LEAVED LEAVES

CAUTION

Avoid contact with skin, eyes, and clothing. Avoid spray mist. Do not reuse container. Destroy when empty.

AVOID INHALING SPRAY MIST

Under no circumstances should this herbicide or any 2,4-D or 2,4,5-T weed killer be used in the vicinity where cotton, tomatoes, grapes, garden crops, ornamentals or other susceptible crops are grown. Extreme care must be exercised to prevent drifting of this material. Do not apply on a windy day. (Coarse sprays are less likely to drift.) Vapors from this product at very high temperatures may cause injury to susceptible plants growing nearby. Do not use spray equipment which has contained this product for other purposes. Do not store near fertilizers, seeds, insecticides, fungicides or foodstuffs. Avoid contamination of water supplies that may be used to spray, irrigate, or water susceptible crops or used for domestic purposes. Do not allow animals to graze on treated turf.

Do not use, pour, spill, or store near heat or open flame. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not apply to forest lands or management areas, around houses or gardens, ditchbanks, water, rights-of-way, pastures or recreational areas.

NOTICE Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

6 Teaspoons = 2 Tablespoons = 1 oz.
12-7
E.P.A. Reg No 2100-241 E.P.A. Est. No. 2560-MS-1

Net Contents 1 Quart

337