

2169-24

March 7, 1975  
EPA REG. NO. 2169-24

**CONTACT SPRAY**—For whiteflies apply Patterson's White Fly Spray at 1 pint per 100 gallons of water or 1 teaspoonful per gallon. Spray upper and lower surfaces of plant foliage to point of runoff. Spray in late afternoon and at night only, when the temperature ranges from 50°F to 72°F. Repeat application weekly for four weeks or longer for control of adults, pupae, and nymphs. Complete spray schedule on poinsettia before blooms form. Where fewer applications are preferred, spray at the rate of 1 qt. per 100 gallons of water or 2 teaspoonfuls per gallon. Repeat as necessary. Apply sprays as mentioned above.

**NON-THERMAL APPLICATORS**—Close off greenhouse area to be treated for whitefly infestations. Direct mist first toward upper corners of the greenhouse, then apply evenly throughout the area. Keep nozzle at least three feet from plant foliage. Apply Patterson's White Fly Spray at the rate of 1½ fl. oz. thoroughly mixed in 3 fl. ozs. of methylene chloride per 10,000 cu. ft. This is equivalent of 1 gram of active ingredient per 1,000 cu. ft. Keep treated area closed off for at least two hours. Do not remain in treated areas. Ventilate the area after treatment is completed. Apply at weekly intervals for at least four weeks. Applications should be made during the late afternoon or at night when the temperature range is 50°F to 72°F.

E.P.A. Reg. No. 2169-24, 373

*patterson's*



Active Ingredients:  
\*†(5-Benzyl-3-furyl)methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate . . . 24.30% w/w  
Related compounds . . . . . 3.30%  
Aromatic petroleum hydrocarbons . . . . . 66.40%  
Inert Ingredients: . . . . . 6.00%  
\*Cis trans isomers ratio: max. 30% (+ or -) cis and min. 70% (+ or -) trans. U.S. Patent No. 3,465,007

**CAUTION: Keep Out of Reach of Children**

See Side Panel for Additional Caution Statements  
Distributed by

**CAUTION: Keep Out of Reach of Children.**  
Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Keep container closed and store at temperatures above 40°F.

This product is toxic to fish and birds. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**FOOD AREAS**—Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation.

**CONTAINER DISPOSAL** Do not reuse container. Crush used container and bury in non cropland soil away from water.

**NOTICE**—Do not store below 40°F. If exposed to temperatures below 40°F warm to 80°F and mix well before using.

**GREENHOUSE PESTS**  
For infestations of whiteflies on begonia, fuchsia, salvia, poinsettia, chrysanthemums, coleus, geraniums, ageratum, petunia, aster, calendula, marigold, roses, azalea and ivy.

**NET CONTENTS ONE QUART**