

DIRECTIONS FOR USE

BAGWORMS — On evergreens and ornamental plants — Mix 3 to 4 teaspoonfuls of 60% Toxaphene per gallon of water. (equal to 3 to 4 pints per 100 gallons). Make sure to add spreader sticker to the solution for best control. Thoroughly spray all foliage when small worms appear the first part of July. Do not spray Red Bud trees since defoliation will occur. If worms are up to ½ to ¾ inches long use the higher rate of application.

CATERPILLARS — On ornamental trees — For control of Walnut caterpillars, tent caterpillars, catalpa worms on catalpa, canker-worms and fall webworms mix 3 to 4 teaspoonfuls per gallon of water (3 to 4 pints per 100 gallons). For caterpillars that produce webbing use spreader sticker to penetrate the web and a sprayer that will produce enough pressure to help the web penetration. Tent caterpillars should be sprayed as the webbing appears. Spring cankerworms on elms and ornamental crabs will require an early spray application on or about May 1st as they appear to eat foliage.

ELM LEAF BEETLE — Elm leaf beetle damage occurs after trees are in full leaf stage therefore spray on or about May 15 to June 1. A second generation will occur within 4 to 5 weeks after the first damage therefore requiring a second spray at the same rate of 5 to 6 teaspoonfuls per gallon. Best control will be accomplished when using spreader sticker in the spray solution.

SOD WEBWORMS — For control of the Blue Grass sod webworm common to the midwest — rake the yard in June and remove the thatch. Spray the area using 2 tablespoons in 5 gallons of water to cover 1000 sq. ft. Make at least 2 more applications at 3 to 4 week intervals to control additional broods that will appear during the warmer weather. Note — The sod webworm has become resistant to Toxaphene in some localities. If this appears to be the case switch your spray to another recommended product.

LIVESTOCK USES

For controlling horn flies on beef cattle, lice, mites (sarcoptic mange/scab mites) on beef cattle, hogs, sheep and goats; and ticks on beef cattle, sheep and goats.

Mixing Rate: Dilute with water and use either as a dip or spray. For a 0.5% toxaphene water spray, mix 3 quarts Toxaphene Concentrate in 100 gallons water, 1 quart in 33 gallons water, 1 pint in 16 gallons or 1 cup (8 oz.) in 8 gallons water.

HORN FLIES, TICKS AND LICE (Beef Cattle, Sheep and Goats): Dip or Spray (When dipping make sure the animal is held in the vat



ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene*) 60.0%

Aromatic Petroleum Derivative Solvent 35.0%

INERT INGREDIENTS: 5.0%

*Containing 67% to 69% Chlcrine.

WARNING:


Keep Out of Reach of Children
See Side Panel for Additional Warnings

NET CONTENTS: 1 QT.

Patterson

E.P.A. Reg. No. 2169-245 5-74

at least
ping.) F
ing or
quarts p
If reinf
Note--D
age. Do
HOGS -
pigs fra
Do not
SARCOF
cattle,
to herd
if neces
6 month
slaughte
BACKRU
dilute
oil, or
cable at
3 week
backrub
location
Note: D
to treat
treatment
NOTE: E
heated d
NOTICE:
handling
label.
WARNIN
Do not
using. D
storage
should r
This pro
other w
streams,
Do not
treated.
disposal
label.
Do not
DO NOT



ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene*)	60.0%
Aromatic Petroleum Derivative Solvent	35.0%

INERT INGREDIENTS: 5.0%

*Containing 67% to 69% Chlcrine.

WARNING:

Keep Out of Reach of Children
See Side Panel for Additional Warnings

NET CONTENTS: 1 QT.

Patterson

E.P.A. Reg. No. 2169-245 5-74

at least two minutes and submerge each animal twice during dipping.) For sheep tick apply during warm weather soon after shearing or before wool is 2 inches long. When spraying use about 2 quarts per beef animal and proportionately less for smaller animals. If reinfestations occur repeat application in 3 weeks for control. Note—Do not apply to dairy animals or animals under 6 months of age. Do not treat animals within 28 days of slaughter.

HOGS — Lice — Spray sows 30 days before farrowing to prevent pigs from being infested. Do not treat pigs under 6 months of age. Do not treat meat hogs within 28 days of slaughter.

SARCOPTIC MANGE (Beef Cattle, Sheep and Goats): Dip or spray cattle, dip sheep and goats, for thorough animal coverage. Apply to herd when mange is found. Repeat treatment in 15 to 21 days if necessary. Note: Do not apply to dairy animals or animals under 6 months of age. Do not treat meat animals within 28 days of slaughter.

BACKRUBBER USE: To control hornflies and aid in reducing lice, dilute 1 part Toxaphene Concentrate with 11 parts kerosene, fuel oil, or diesel oil. Apply this diluted solution to canvas or burlap cable at the rate of one gallon to 20 feet of cable. Repeat at 2 to 3 week intervals or as necessary to keep cable dampened. Locate backrubber to give beef cattle and pigs free access to it. Best locations are usually near watering places, salt licks or feed racks. Note: Do not allow calves under 6 months of age, or dairy animals to treat themselves. Do not permit meat animals access to this treatment within 28 days of slaughter.

NOTE: Do not treat animals during cold, stormy weather and overheated or sick animals.

NOTICE: Buyer assumes all risk and liability for use, storage and/or handling of this material not in accordance with the terms of this label.

WARNING: May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed and food products. Not for use or storage in or around the home. This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

DIRECTIONS FOR USE

Evergreens and ornamental plants - Mix 3 to 9% Toxaphene per gallon of water. Equal to 3 gallons. Make sure to add spreader sticker to get control. Thoroughly spray all foliage when the first part of July. Do not spray Red Bud until worms are up to 1/2 to 3/4 inches rate of application.

On ornamental trees - For control of Walnut caterpillars, catalpa worms on catalpa, cankerworms mix 3 to 4 teaspoonfuls per gallon of water per 100 gallons. For caterpillars that produce webbing use a spreader sticker to penetrate the web and a sprayer with enough pressure to help the web penetration. Trees should be sprayed as the webbing appears. Spring and ornamental crabs will require an early application or about May 1st as they appear to eat


Elm leaf beetle damage occurs after trees are therefore spray on or about May 15 to June 1. Damage will occur within 4 to 5 weeks after the first application requiring a second spray at the same rate of application per gallon. Be finished when using spreader sticker in the spray

For control of the Blue Grass sod webworm - rake the yard in June and remove the sod using 2 tablespoons in 5 gallons of water. Make at least 2 more applications at 3 to 4 week intervals. Control additional broods that will appear during the season. Note - The sod webworm has become resistant in some localities. If this appears to be the case spray to another recommended product

LIVESTOCK USES

On flies on beef cattle, lice, mites, sarcoptic mange on beef cattle, hogs, sheep and goats, and ticks on sheep and goats. Use either as a dip or spray. For one water spray, mix 3 quarts Toxaphene Concentrate in 33 gallons water. 1 pint in 8 gallons water.

AND LICE - Beef Cattle, Sheep and Goats: Dip or spray make sure the animal is held in the vat



ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene*) 60.0%
Aromatic Petroleum Derivative Solvent 35.0%

INERT INGREDIENTS:

5.0%

*Containing 67% to 69% Chlorine.

WARNING:

Keep Out of Reach of Children
See Side Panel for Additional Warnings

NET CONTENTS: 1 QT.

at least two minutes and submerge feet during dipping. For sheep tick apply during warm weather or before wool is 2 inches long. When 3 quarts per beef animal and proportionately less for hogs. If reinfestations occur repeat application in 10 days. Note-Do not apply to dairy animals or animals under 6 months of age. Do not treat animals within 28 days of slaughter.

HOGS - Lice - Spray sows 30 days before pigs from being infested. Do not treat piglets. Do not treat meat hogs within 28 days of slaughter.

SARCOPTIC MANGE - Beef Cattle, Sheep and Goats - Dip sheep and goats, for thorough application to herd when mange is found. Repeat treatment if necessary. Note: Do not apply to dairy animals under 6 months of age. Do not treat meat animals within 28 days of slaughter.

BACKRUBBER USE: To control hornflies and mosquitoes dilute 1 part Toxaphene Concentrate with 11 parts of kerosene, diesel oil, or diesel fuel. Apply this diluted solution to the backrubber at the rate of one gallon to 20 feet of backrubber at 3 week intervals or as necessary to keep backrubber to give beef cattle and hogs freedom of movement. Locations are usually near watering places, salt licks, etc. Note: Do not allow calves under 6 months of age to treat themselves. Do not permit animals to be treated within 28 days of slaughter.

NOTE: Do not treat animals during cold, storm, or when heated or sick animals.

NOTICE: Buyer assumes all risk and liability for handling of this material not in accordance with label.

WARNING: May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Use with care. Do not store near feed and food products. Do not use in or around the home. This product should not be applied when bees are actively flying. This product is toxic to fish, birds, and other wildlife in treated areas may be killed. Do not apply when weather conditions favor runoff. Do not apply when weather conditions favor runoff. Do not contaminate water by clean disposal of wastes. Apply this product only as directed on label.

Do not reuse container. Destroy when empty. DO NOT USE POUR, SPILL OR STORE IN A PLACE