

**DIRECTIONS FOR USE**

KERB 50W is a wettable powder and should be premixed into a thin paste with water before adding to the spray tank. Use return-flow or mechanical agitation to keep the product in suspension. If a hose-end lawn sprayer or a pressure type garden sprayer is used, agitate the spray solution by shaking the mixture frequently when applying. When spraying small lawn areas apply at least 2 gallons of spray solution per 1000 sq. ft. On average basis use 20 to 50 gallons per acre.

KERB acts mainly through root absorption. For this reason if no rain occurs after application within a day, lightly irrigate the area to move the chemical into the soil. The chemical works slowly on seedling and mature grasses. Excessive irrigation or heavy rains will wash and leach the chemical from the area of application.

**POA ANNUA CONTROL:** Bermudagrass golf courses, Lawns and Playing fields. Pre-emergence and early postemergence control—Use 1 to 2 lbs. KERB 50W per acre (2 1/2 to 4 2/3 tablespoons per 2 to 4 gallons of water per 1000 sq. ft.). In areas of heavy growth use the higher rates of KERB and water to get best control and longer residual control. Apply in the fall or late winter when POA ANNUA normally begins to germinate or is in the one to two leaf stage.

Pre-Emergence—Use 1 1/2 to 2 lbs. KERB 50W per acre (3 1/2 to 4 2/3 tablespoons per 2 to 4 gallons of water per 1000 sq. ft.) when POA ANNUA is in the early tillering stage (multi-grass shoots from plant) to the seed heading stage. If seed heads have already formed increase rate to 2 to 3 pounds per acre (4 2/3 to 7 tablespoons per 1000 sq. ft.).

**PROTECTIVE CLOTHING:** Use the following items of clothing when mixing or applying this product: long-sleeved shirts and long pants, preferably one-piece (coveralls), hat with brim, heavy-duty fabric or rubber work gloves. Use heavy-duty leather or rubber boots when applying with hand spray equipment.

2169-248

**CAUTION!**  
Keep out of reach of children.  
See Additional Caution Statements on Side Panel  
Distributed by

NOTE—Dichondra, bluegrass, annual and perennial bentgrass are sensitive to pre and post-emergence control. Do not overseed with susceptible cool season grasses. Avoid spraying areas on fairways or approaches to greens that have been overseeded with cool season grasses. KERB 50W is not recommended on greens as adequate control is not yet available.

**PRECAUTIONARY STATEMENTS:** Hazardous to Humans and Domestic Animals. Harmful if swallowed. Avoid skin and eye contact. Wash with soap and water. In case of eye contact wash thoroughly with water. Do not breathe the dust or spray mist.

**ENVIRONMENTAL HAZARD:** STORAGE AND DISPOSAL. Apply only as specified on this label. Do not reuse for any other purpose. In strict accordance with label directions bury in a safe place in non-crop lands away from buildings.

**NOTICE:** Buyer assumes all risk and liability of the material for its use in accordance with the terms of the label.

U.S. Patents 2,842,000 and 2,842,001  
E.P.A. Reg. No. 2169-248  
Kerb Reg. T.M. Rohm & Haas Co.

**RESTRICTED USE PESTICIDE** For retail sale to and application only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators Certification

**DIRECTIONS FOR USE**

KERB 50W is a wettable powder and should be premixed into a thin paste with water before adding to the spray tank. Use return-flow or mechanical agitation to keep the product in suspension. If a hose-end lawn sprayer or a pressure type garden sprayer is used, agitate the spray solution by shaking the mixture frequently when applying. When spraying small lawn areas apply at least 2 gallons of spray solution per 1000 sq. ft. On average basis use 20 to 50 gallons per acre.

KERB acts mainly through root absorption. For this reason if no rain occurs after application within a day, lightly irrigate the area to move the chemical into the soil. The chemical works slowly on seedling and mature grasses. Excessive irrigation or heavy rains will wash and leach the chemical from the area of application.

**POA ANNUA CONTROL:** Bermudagrass golf courses, Lawns and Playing fields. Pre-emergence and early postemergence control—Use 1 to 2 lbs. KERB 50W per acre (2 1/2 to 4 2/3 tablespoons per 2 to 4 gallons of water per 1000 sq. ft.). In areas of heavy growth use the higher rates of KERB and water to get best control and longer residual control. Apply in the fall or late winter when POA ANNUA normally begins to germinate or is in the one to two leaf stage.

Pre-Emergence—Use 1 1/2 to 2 lbs. KERB 50W per acre (3 1/2 to 4 2/3 tablespoons per 2 to 4 gallons of water per 1000 sq. ft.) when POA ANNUA is in the early tillering stage (multi-grass shoots from plant) to the seed heading stage. If seed heads have already formed increase rate to 2 to 3 pounds per acre (4 2/3 to 7 tablespoons per 1000 sq. ft.).

**PROTECTIVE CLOTHING:** Use the following items of clothing when mixing or applying this product: long-sleeved shirts and long pants, preferably one-piece (coveralls), hat with brim, heavy-duty fabric or rubber work gloves. Use heavy-duty leather or rubber boots when applying with hand spray equipment.

pattersons

**KERB™**  
50-W  
A selective herbicide for Poa Annuua Control in Bermudagrass

**CAUTION!**  
Keep out of reach of children.  
See Additional Caution Statements on Side Panel  
Distributed by

NOTE—Dichondra, bluegrass, annual and perennial bentgrass are sensitive to pre and post-emergence control. Do not overseed with susceptible cool season grasses. Avoid spraying areas on fairways or approaches to greens that have been overseeded with cool season grasses. KERB 50W is not recommended on greens as adequate control is not yet available.

**CAUTION:** Harmful if swallowed. Avoid skin and eye contact. Wash with soap and water. In case of eye contact wash thoroughly with water. Do not breathe the dust or spray mist.

**ENVIRONMENTAL HAZARD:** STORAGE AND DISPOSAL. Apply only as specified on this label. Do not reuse for any other purpose. In strict accordance with label directions bury in a safe place in non-crop lands away from buildings.

**NOTICE:** Buyer assumes all risk and liability of the material for its use in accordance with the terms of the label.

U.S. Patents 3,534,096 and 3,534,097  
E.P.A. Reg. No. 2169-248  
Kerb Reg. T.M. Rohm & Haas Co.

**NET CONTENTS: 6 OZ**

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

**INSTRUCTIONS FOR USE**  
 should be premixed into a thin paste with water. Use return-flow or mechanical agitation to mix. Use lawn sprayer or a pressure type garden sprayer by shaking the mixture frequently when areas apply at least 2 gallons of spray solution per acre. For this reason if no rain occurs after the area to move the chemical into the root feeding and mature grasses. Excessive irrigation the chemical from the area of application. Treat golf courses, lawns and playing fields. Control—Use 1 to 2 lbs. KERB 50W per acre (4 gallons of water per 1000 sq. ft.). In areas of KERB and water to get best control and longer life winter when POA ANNUA normally begins seed stage. KERB 50W per acre (1 1/2 to 2 1/2 tablespoons per acre) when POA ANNUA is in the early tillering stage. In the seed heading stage, if seed heads have 3 pounds per acre (4 1/2 to 7 tablespoons per

Following items of clothing when mixing or applying and long pants, preferably one-piece (overalls), rubber work gloves. Use heavy-duty leather or spray equipment.

2

Use only by Certified Applicators

**NOTE**—Dichloro, Monogras, annual and perennial ryegrasses, fescues, Zoysia and bentgrass are sensitive to pre and post-emergence application of KERB. Do not overseed with susceptible cool season grasses within 90 days of treatment. Avoid spraying areas on fairways or approaches that drain into bentgrass greens or areas that have been overseeded with cool season grasses. KERB 50W is not recommended on greens as adequate information for this application is not yet available.

**PRECAUTIONARY STATEMENTS**  
 Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed. Avoid skin and eye contact. In case of skin contact wash with soap and water. In case of eye contact wash with large amounts of water. Do not breathe the dust or spray mist.

**ENVIRONMENTAL HAZARDS**  
 STORAGE AND DISPOSAL

Apply only as specified on this label. Do not reuse container. Dispose of by perforating and either burning in strict accordance with air pollution control laws or by burying in a safe place in non-crop lands away from water supplies.

**NOTICE:** Buyer assumes all risk and liability of use, storage, and/or handling of this material and in accordance with the terms of this label.

U.S. Patents 2,894,898 and 3,640,899

ACTIVE INGREDIENTS:	3,5-dichloro-N (1,1-dimethyl-2-propenyl) benzamide	50.0%
INERT INGREDIENTS:	Total	50.0%
	Total	100.0%

E.P.A. Reg. No. 249-548 E.P.A. Est. 2169-Ka-1  
 Herb Res. T. M. Rohm & Haas Co., Philadelphia, Pa.


2169-248

**CAUTION!**  
 Keep out of reach of children.  
 See Additional Caution Statements on Side Panel  
 Distributed by

**RESTRICTED USE PESTICIDE** For retail sale to and application only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators Certification

**INSTRUCTIONS FOR USE**  
 should be premixed into a thin paste with water. Use return-flow or mechanical agitation to mix. Use lawn sprayer or a pressure type garden sprayer by shaking the mixture frequently when areas apply at least 2 gallons of spray solution per acre. For this reason if no rain occurs after the area to move the chemical into the root feeding and mature grasses. Excessive irrigation the chemical from the area of application. Treat golf courses, lawns and playing fields. Control—Use 1 to 2 lbs. KERB 50W per acre (4 gallons of water per 1000 sq. ft.). In areas of KERB and water to get best control and longer life winter when POA ANNUA normally begins seed stage. KERB 50W per acre (1 1/2 to 2 1/2 tablespoons per acre) when POA ANNUA is in the early tillering stage. In the seed heading stage, if seed heads have 3 pounds per acre (4 1/2 to 7 tablespoons per

Following items of clothing when mixing or applying and long pants, preferably one-piece (overalls), rubber work gloves. Use heavy-duty leather or spray equipment.



**CAUTION!**  
 Keep out of reach of children.  
 See Additional Caution Statements on Side Panel  
 Distributed by

**NOTE**—Dichloro, Monogras, annual and perennial ryegrasses, fescues, Zoysia and bentgrass are sensitive to pre and post-emergence application of KERB. Do not overseed with susceptible cool season grasses within 90 days of treatment. Avoid spraying areas on fairways or approaches that drain into bentgrass greens or areas that have been overseeded with cool season grasses. KERB 50W is not recommended on greens as adequate information for this application is not yet available.

**CAUTION:** Harmful if swallowed. Avoid skin and eye contact. In case of skin contact wash with soap and water. In case of eye contact wash with large amounts of water. Do not breathe the dust or spray mist.

Apply only as specified on this label. Do not reuse container. Dispose of by perforating and either burning in strict accordance with air pollution control laws or by burying in a safe place in non-crop lands away from water supplies.

**NOTICE:** Buyer assumes all risk and liability of use, storage, and/or handling of this material and in accordance with the terms of this label.

U.S. Patents 3,534,898 and 3,640,899

ACTIVE INGREDIENTS:	3,5-dichloro-N (1,1-dimethyl-2-propenyl) benzamide	50.0%
INERT INGREDIENTS:	Total	50.0%
	Total	100.0%

E.P.A. Reg. No. 249-548 E.P.A. Est. 2169-Ka-1  
 Herb Res. T. M. Rohm & Haas Co., Philadelphia, Pa.

**NET CONTENTS: 6 OUNCES**

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE